

2004 HIGHER SCHOOL CERTIFICATE SOLUTIONS GENERAL MATHEMATICS

SECTION I

SUMMARY

1. B 2. B 3. D 4. A 5. D 6. C 7. D 8. B 9. C 10. C 11. B 12. C 13. C 14. A 15. A 16. D 17. A 18. C 19. B 20. A 21. D 22. A

1. (B) $25\% = \frac{25}{100} = \frac{1}{4}$

2. (B) Using the points (0, 10) and (4, 30):

$$\begin{aligned} \text{Gradient} &= \frac{\text{vertical rise}}{\text{horizontal run}} \\ &= \frac{30-10}{4-0} \\ &= \frac{20}{4} \\ &= 5. \end{aligned}$$

3. (D) $K = Ft^3$
 $= 5 \times (0.715)^3$
 $= 1.827 \ 629 \ 375 \dots$
 ≈ 1.83 (3 significant figures).

4. (A) Real estate agents receive a commission for selling property.

5. (D) $\tan 20^\circ = \frac{\text{opposite}}{\text{adjacent}} = \frac{c}{a}$

6. (C) Range = highest score - lowest score
 $= 12 - 1$
 $= 11$

7. (D) 1, 3, 3, 3, 4, 5, 7, 7, 12

Median is the middle score for an odd set of scores. Median = 4.
 Mode is the score with the highest frequency. Mode = 3.

8. (B) Sector angle for school X = 100° .
 Number of prizes won by school X
 $= \frac{100 \times 116}{360}$
 $= 32.2222 \dots$
 ≈ 32

\therefore School X won 32 prizes (number of prizes must be a whole number).

9. (C) Area = $\frac{1}{2} ab \sin C$

$$\begin{aligned} &= \frac{1}{2} \times 30 \times 20 \times \sin 35^\circ \\ &= 172.072 \ 930 \ 9 \dots \\ &\approx 172 \text{ m}^2 \text{ (nearest m}^2\text{)}. \end{aligned}$$

10. (C) Tax payable
 $= \$8310 + 0.4 \times (\$47 \ 000 - \$45 \ 000)$
 $= \$9110.00$

11. (B) $d = 6t^2$
 $\frac{2400}{6} = \frac{6t^2}{6}$
 $400 = t^2$
 $t = \pm \sqrt{400}$
 $t = \pm 20$

$\therefore t = 20$ is a possible value.

12. (C) Interquartile range
 $=$ upper quartile - lower quartile
 $= 11 - 8$
 $= 3$

13. (C) Kath breathes $10 \times 100 \text{ L} = 1000 \text{ L}$.
 Jim breathes $10 \times 6 \text{ L} = 60 \text{ L}$.
 Difference = $1000 \text{ L} - 60 \text{ L} = 940 \text{ L}$.

14. (A) Financial expectation
 $= \frac{1}{10} \times \$20 + \frac{1}{2} \times \$1 + \frac{2}{5} \times (-\$2)$
 $= \$1.70$

15. (A) Shaded area
 $=$ area of ellipse - area of rectangle.
 For the ellipse: For the rectangle:
 $a = \frac{1}{2} \times 60 = 30$ $L = 24$
 $b = \frac{1}{2} \times 40 = 20$ $B = 10$
 \therefore Area = $\pi ab - LB$
 $= \pi \times 30 \times 20 - 24 \times 10$
 $= 1644.955 \ 592 \dots$
 $\approx 1645 \text{ mm}^2$ (nearest mm^2).

16. (D) Both linear equations are of the form $y = mx + b$ where $m =$ gradient, $b =$ y-intercept.

\therefore The graph of $y = 2x - 5$ has a gradient of 2 and a y-intercept of -5.
 \therefore The graph of $y = x + 6$ has a gradient of 1 and a y-intercept of 6.
 (D) is the only diagram that shows 2 lines, both having positive gradient, one line with a positive y-intercept and the other line with a negative y-intercept.

METHOD 1

110% of original price = \$880.
 \therefore 10% = $\frac{\$880}{11}$
 \therefore GST refund = \$80.

METHOD 2

110% of original price = \$880.
 \therefore 1% = $\frac{\$880}{110}$
 \therefore 10% = \$80
 \therefore GST refund = \$80.

18. (C)	1	2	3	4	5	6
1	•	•	•	•	•	• (1, 6)
2	•	•	•	•	•	• (2, 6)
3	•	•	•	•	•	• (3, 6)
4	•	•	•	•	•	• (4, 6)
5	•	•	•	•	•	• (5, 6)
6	• (6, 1)	• (6, 2)	• (6, 3)	• (6, 4)	• (6, 5)	•

There are 36 possible outcomes when 2 dice are rolled.

$P(\text{only one dice shows } 6) = \frac{10}{36} = \frac{5}{18}$

19. (B) $A = P(1+r)^n$ where $A =$ final amount
 $P = 250$
 $r = 0.05\%$
 $= \frac{0.05}{100}$ per day
 $n = 30$ days.

$$\begin{aligned} A &= 250 \times \left(1 + \frac{0.05}{100}\right)^{30} \\ &= 253.777 \ 314 \ 8 \dots \\ \text{Final value} &= \$253.78 \text{ (to nearest cent).} \\ \therefore \text{Interest charged} &= \$253.78 - \$250 \\ &= \$3.78. \end{aligned}$$

METHOD 1

Equivalent normal hours worked
 $= 20 + 4 \times 1.5$
 $= 26$.
 So hourly rate = $\frac{\$291.20 + 26}{26} = \11.20 .

METHOD 2

Let hourly rate be h .
 $20 \times h + 4 \times 1.5 \times h = 291.20$
 $20h + 6h = 291.20$
 $26h = 291.20$
 $h = \frac{291.20}{26}$
 $= 11.2$.

\therefore Hourly rate is \$11.20.

21. (D) Relationship is of the form $t = \frac{k}{n}$, where k is a constant.

A graph of this relationship would be a hyperbola. Note that a hyperbola is a curve that approaches each axis but has no point on either axis.

22. (A) One Australian dollar = 0.62 euros
 \therefore to convert Australian dollars to euros multiply by 0.62.
 \therefore Number of euros in 25 Australian dollars = 25×0.62
 $= 15.5$ euros.

One Vistabella dollar = 1.44 euros
 \therefore to convert euros to Vistabella dollars divide by 1.44.

\therefore Number of Vistabella dollars in 15.5 euros = $\frac{15.5}{1.44}$
 $= 10.763 \ 888 \ 89 \dots$
 $= \$10.76$.

SECTION II

Question 23


(a) (i) Garden area
 $=$ area of $\triangle ABC$ + area of $\triangle ADC$
 $= \frac{1}{2} \times 10 \times 5.1 + \frac{1}{2} \times 10 \times 6.3$
 $= 57 \text{ m}^2$.

(ii) $V = Ah$ where $A = 57 \text{ m}^2$, $h = 5 \text{ cm} = 0.05 \text{ m}$
 \therefore Volume of straw = 57×0.05
 $= 2.85 \text{ m}^3$.

(iii) Number of bags = $\frac{2.85}{0.25}$
 $= 11.4$
 ≈ 12 (rounded up).
 \therefore Carnel needs 12 bags of straw.

(iv)

$AB^2 = 5.1^2 + 6.0^2$ (Pythagoras' theorem)
 $= 62.01$.



$$AB = \sqrt{62.01}$$

$$= 7.874\ 642\ 849 \dots$$

$$\approx 8\text{ m.}$$

\therefore Length of the fence = 8 m (to the nearest metre).

(b) (i) Cost of Novirus = $x + 115$.

(ii) The equation is $x + x + 115 = 415$
 $2x + 115 = 415$
 $2x = 300$
 $x = 150$.

The cost of Funmaths is \$150.00.

By similar triangles

$$\frac{h}{2.7} = \frac{5}{2}$$

$$h = \frac{5 \times 2.7}{2}$$

$$= 6.75.$$

\therefore The height of the tree is 6.75 metres.

Question 24


(a) (i) The mean 3 pm wind speed for September is 15 km/h.

(ii) The month with the lowest 3 pm wind speed is February.

(iii) The highest number of raindays occur in the three-month period January, February, March.

(iv) When the wind speed is high, as in July, August and September, the number of raindays is low. When the wind speed is low, as in February and March, the number of raindays is high.

(b) (i)




Let C lie due east of A.

$$\angle PAQ = \angle PAC + \angle CAQ$$


$$= 90^\circ + 45^\circ$$

$$= 135^\circ.$$

(ii)



(iii)



Using the cosine rule

$$\cos \angle PAB = \frac{P^2 + B^2 - A^2}{2PB}$$

$$= \frac{31^2 + 28^2 - 35^2}{2 \times 31 \times 28}$$


$$= 0.299\ 539\ 17 \dots$$

$$\therefore \angle PAB = 72.570\ 073\ 25 \dots$$

$$\approx 73^\circ \text{ (correct to nearest degree).}$$

(c) (i) Roberto's score is on the line between 180 and 190 (below the shaded area). It is a score between 1 and 2 standard deviations above the mean, so his z-score is a number between 1 and 2.

(ii) METHOD 1




For a normal distribution:

68% of scores lie within 1 standard deviation of the mean.
 95% of scores lie within 2 standard deviations of the mean.

$$\therefore \text{shaded area} = \frac{1}{2} \times (95\% - 68\%)$$

$$= 13.5\%.$$

METHOD 2



For a normal distribution:

68% of scores lie within 1 standard deviation of the mean.
 \therefore 34% of scores lie within the mean and 1 standard deviation above the mean.
 95% of scores lie within 2 standard deviations of the mean.
 \therefore 47.5% of scores lie within the mean and 2 standard deviations above the mean.

$$\therefore \text{shaded area} = 47\frac{1}{2}\% - 34\%$$

$$= 13\frac{1}{2}\%.$$

Question 25

(a) (i) METHOD 1

$$2003 \text{ tax deduction} = 40\% \text{ of } \$6500$$

$$= 0.4 \times \$6500$$

$$= \$2600.$$

METHOD 2

$$S = V_0(1+r)^n$$

$$S = 6500$$

$$r = 40\% = 0.40$$

$$n = 1$$

The value of the computer after one year is

$$S = 6500 \times (1 - 0.40)^1$$

$$= \$3900.$$

$$\therefore \text{The tax deduction} = \$6500 - \$3900$$

$$= \$2600.$$

(ii) After 3 years, $S = 6500 \times (1 - 0.40)^3$
 $= \$1404.$

\therefore The value of her computer at the start of the 2006 financial year is \$1404.

(b) (i) METHOD 1

The arrangements are: CSV, CVS
 VSC, VCS
 SVC, SCV

\therefore There are 6 different ways of arranging the tubs in a row.

METHOD 2

There are 3 choices for 1st tub, 2 for 2nd tub, 1 for 3rd tub.
 Number of choices = $3 \times 2 \times 1$
 $= 6.$

\therefore There are 6 different ways of arranging the tubs in a row.

(ii) METHOD 1


$$P(\text{choosing chocolate on one day}) = \frac{1}{3}.$$

$$P(\text{choosing chocolate on both days})$$

$$= \frac{1}{3} \times \frac{1}{3}$$

$$= \frac{1}{9}.$$

METHOD 2



$$\therefore P(\text{choosing chocolate on both days}) = \frac{1}{9}.$$

(iii) METHOD 1

The two flavours in this case must be different. There are 3 choices for the first flavour, 2 choices for the second flavour.

$$P(\text{selecting chocolate as first flavour}) = \frac{1}{3}.$$

$$P(\text{selecting strawberry as second flavour}) = \frac{1}{2}.$$


$$\therefore P(\text{selecting chocolate first then strawberry})$$

$$= \frac{1}{3} \times \frac{1}{2}$$

$$= \frac{1}{6}.$$

METHOD 2

The two flavours in this case must be different.



$$\therefore P(\text{chocolate first, then strawberry}) = \frac{1}{6}.$$

(c) (i)	Test indicated a lie	Test did not indicate a lie	Total
People who lied	40	10	50
People who did not lie	20	130	150

(ii) Number of people tested where lie-detector test was accurate = $40 + 130$ from table = 170.

(iii) Percentage of people tested where the test was accurate = $\frac{170}{200} \times 100\%$ = 85%.

Note: Denominator is 200 as required percentage is based on all the people tested.

(iv) Note: In this case only referring to the people who did not lie, so initial denominator will be 150.

$$P(\text{test indicated a lie for a person who did not lie}) = \frac{20}{150} = \frac{2}{15}$$


Question 26

(a) (i) Percentage increase = $\frac{14}{100} \times 100\%$ = 14%.

(ii) Given that $n = 100(1.14)^t$.
When $t = 15$, $n = 100(1.14)^{15}$
= 713.793 797 8 ...
= 714.

\therefore There are 714 bacteria after 15 hours.

(iii) Graph of $n = 100(1.14)^t$.



(iv) From the graph, $n = 300$ when $t \approx 8$ so the population is 300 after 8 hours.

An alternative method is to use guess and check to solve $300 = 100(1.14)^t$.

(b) (i) 131°E

Sorong 1°S

Darwin 12°S

Difference in latitude = $12^\circ - 1^\circ$ = 11° .

METHOD 1

An angular distance of 1° on a great circle equals 60 nautical miles.

$$\begin{aligned} \text{Distance} &= 11 \times 60 \text{ nautical miles} \\ &= 11 \times 60 \times 1.852 \text{ km} \\ &= 1222.32 \text{ km} \\ &\approx 1200 \text{ km}. \end{aligned}$$

METHOD 2

The arc length formula can be used.

$$\begin{aligned} \ell &= \frac{\theta}{360} 2\pi r \text{ where } \theta = 11^\circ \\ r &= 6400 \\ \text{Distance} &= \frac{11}{360} \times 2\pi \times 6400 \\ &= 1228.711 793 \dots \\ &\approx 1200 \text{ km}. \end{aligned}$$

METHOD 1

Assuming the tourists travel in a straight line, the distance is approximately 1200 km, or $1200 \div 1.852 \approx 647.95$ nautical miles. At a speed of 15 nautical miles per hour (ie. 15 knots) this would take $647.95 \div 15 \approx 43.2$ hours

$$\left(\text{since time} = \frac{\text{distance}}{\text{speed}} \right).$$

So they would complete the trip in less than 48 hours.

METHOD 2

Distance = 11×60

= 660 nautical miles.

Using speed = $\frac{\text{distance}}{\text{time}}$, or the

$$\text{diagram } \frac{D}{S} = T,$$

time taken = $\frac{660}{15}$

= 44 hours.

So they would complete the trip in less than 48 hours.

METHOD 3

At 15 knots = 15 nautical miles per hour.

Distance travelled in 48 hours

$$= 15 \times 48$$

$$= 720 \text{ nautical miles}$$

$$= 720 \times 1.852 \text{ km}$$

$$= 1333.44 \text{ km}.$$

So they would be able to travel the distance of 1200 km in less than 48 hours.

Question 27

(a) (i) 150 000 = 150×1000

From the table:

$$\text{Monthly repayments} = 7.75 \times 150 = \$1162.50.$$

$$(ii) \text{ Total repayments} = \$1162.50 \times 20 \times 12 = \$279\,000.$$

$$\text{Total interest repaid} = \$279\,000 - \$150\,000 = \$129\,000.$$

METHOD 1

Interest paid if loan is paid after 15 years

$$\text{is: Interest} = \$242\,730 - \$150\,000 = \$92\,730.$$

$$\therefore \text{Savings in interest is: Difference in interest} = \$129\,000 - \$92\,730 = \$36\,270.$$

METHOD 2

The interest is the difference between the total payments.

$$\text{So difference in interest} = \$279\,000 - \$242\,730 = \$36\,270.$$

METHOD 1

David's Saturday pay = $6 \times 1.5 \times \$18$ = \$162.

Number of hours needed to work at the

$$\text{weekday rate } (\$18/h) = \$162 \div 18 = 9 \text{ hours}.$$

METHOD 2

On a Saturday David works 6 hours

time-and-a-half.

$$\text{Equivalent weekday hours} = 6 \times 1.5 = 9 \text{ hours}.$$

(c) The future value of \$200 per month at 6% pa compounded monthly over 4 years is found using the 'Future value of an annuity' formula.

In this case, $M = \$200$

$$r = \frac{0.06}{12} = 0.005 \text{ per month}$$

$$n = 12 \times 4 = 48 \text{ months}.$$

$$A = M \left[\frac{(1+r)^n - 1}{r} \right]$$

$$= \$200 \left[\frac{(1+0.005)^{48} - 1}{0.005} \right]$$

$$= \$10\,819.57.$$

\therefore Sanjeev will reach his goal of \$10 500 with \$319.57 in excess.

Question 28

(a) (i) In this case, $R = \frac{w}{h^2}$.

Substitute $w = 72$, $h = 1.50$ (in metres).

$$R = \frac{72}{(1.5)^2}$$

$$= 32.$$

Fred's health rating is 32.

(ii) Substitute $R = 25$, $h = 1.60$ (in metres)

$$25 = \frac{w}{(1.6)^2}$$

$$w = 25 \times (1.6)^2$$

$$= 64 \text{ kg}.$$

Difference in weights = $72 - 64 = 8 \text{ kg}.$

ie. Fred needs to lose 8 kg.

(i) Relationship is of the form $C = kb^3$, where k is a constant.

(ii) Substitute $C = 50$, $b = 10$.

$$50 = k \times 10^3$$

$$50 = 1000k$$

$$k = \frac{50}{1000}$$

$$\therefore k = 0.05.$$

METHOD 1

$$C = 0.05b^3$$

Replace b with $2b$ in equation from (b)(i).

$$C = k(2b)^3$$

$$= k \times 8b^3$$

$$= 8kb^3.$$

\therefore The cost is 8 times bigger, so Felicity is incorrect.

METHOD 2

$$C = 0.05b^3.$$

$$\text{Substitute } b = 10: C = 0.05 \times 10^3 = 50.$$

$$\text{Substitute } b = 20 \text{ (since the base is doubled):}$$

$$C = 0.05 \times (20)^3 = 400.$$

When the base is doubled the cost is 8 times more, so Felicity is incorrect.

- (c) (i) Median height of middle section = 170.
Median weight of middle section = 55.

\therefore Coordinates of C are (170, 55).

- (ii) Jill needs to draw a line parallel to AB which is a third of the distance towards C .

- (iii) (1) Weight in kg = $\frac{2}{3}$ (height in cm) - 50.

$$\text{Substitute weight} = 75.$$

$$75 = \frac{2}{3} (\text{height in cm}) - 50$$

$$125 = \frac{2}{3} \text{ height in cm}$$

$$\text{height in cm} = \frac{125}{2} \times 3$$

$$= 187.5.$$

\therefore The height is predicted to be 187.5 cm.

- (2) Suggested possible answers are:

- the sample used is small (only 9 people)

OR

- tall people are not necessarily heavier

OR

- the model cannot be used outside the range of the sample

OR

- the accuracy with which the graph is drawn.

END OF GENERAL MATHEMATICS SOLUTIONS