

# NEWCASTLE GRAMMAR SCHOOL

Student Number: \_\_\_\_\_



## 2011 TRIAL HIGHER SCHOOL CERTIFICATE EXAMINATION

# Mathematics Extension 1

*Examination Date: Friday 19th August*

Examiner: Mr. M. Brain

### General Instructions

- Reading Time - 5 minutes
- Working Time - 2 hours
- Write using a blue or black pen
- Write your student number on every booklet
- Board-approved calculators may be used
- A table of standard integrals is provided in this paper
- All necessary working should be shown in every question
- Each question attempted is to be returned in a separate Writing Booklet clearly marked Questions 1 etc.
- If required, additional booklets may be requested

### Total marks - 84

- Attempt Questions 1- 7
- All questions are of equal value

**Question 1** (Start a new booklet)

**Marks**

- a) If  $y = (\tan^{-1} x)^2$  find  $\frac{dy}{dx}$  **2**
- b) Find the value of  $\sum_{n=2}^5 {}^n C_2$  **2**
- c) Solve  $\frac{2x-3}{x-2} \geq 1$  **4**
- d) Find  $\lim_{x \rightarrow 0} \frac{\sin 3x}{x}$  **1**
- e) Evaluate  $\int_0^3 \frac{x}{\sqrt{1+x}} dx$  using the substitution  $x = u^2 - 1$ , where  $u > 0$  **3**

**Question 2** (Start a new booklet)

**Marks**

- a) Find the size of the acute angle between the two lines with equations  $x + 2y + 1 = 0$  and  $2x - 3y + 6 = 0$ , correct to the nearest degree **4**
- b) A spherical balloon is expanding so that its volume,  $V \text{ cm}^3$ , increases at a constant rate of  $72 \text{ cm}^3$  per second. What is the rate of increase of the surface area when the radius is 12 cm? **4**
- c) Find **4**
- (i)  $\int \sin^2 x \, dx$
- (ii)  $\int \frac{dx}{\sqrt{9 + 4x^2}}$

**Question 3** (Start a new booklet)

**Marks**

- a) The rate of growth of a bacteria colony is proportional to the excess of the colony's population over 5000 and is given by

**5**

$$\frac{dN}{dt} = k(N - 5000)$$

- (i) Show that  $N = 5000 + Ae^{kt}$  is a solution to the differential equation above
- (ii) Given that the initial population was 15000 and had risen to 20000 after 2 days find the value of  $A$  and  $k$
- (iii) Hence calculate the expected population after 7 days

- b) The polynomial  $P(x) = x^5 + mx^3 + nx$  has a remainder of 5 when divided by  $(x - 2)$ , where  $m$  and  $n$  are constants.

**3**

- (i) Prove that  $P(x)$  is an odd function
- (ii) Hence find the remainder when  $P(x)$  is divided by  $(x + 2)$

- c) If  $\alpha, \beta$  and  $\gamma$  are the roots of the equation  $x^3 - 2x^2 + 3x + 7 = 0$  find the value of  $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$


**4**

**Question 4** (Start a new booklet)

**Marks**

- a)  $ABC$  is a triangle inscribed in a circle.  $M$  is a point on the tangent to the circle at  $B$  and  $N$  is a point on  $AC$  produced so that  $MN$  is parallel to  $BA$

**4**



Copy the diagram into your answer booklet

- (i) State why  $\angle MBC = \angle BAC$
- (ii) Prove that  $MNCB$  is a cyclic quadrilateral
- b) Prove by mathematical induction that, for all integers  $n \geq 1$

**4**

$$1 \times 2^0 + 2 \times 2^1 + 3 \times 2^2 + \dots + n \times 2^{n-1} = 1 + (n-1)2^n$$

- c) (i) Prove that the equation  $\frac{\log_e x}{x} + 2 = 0$  has a solution between  $x = 0.4$  and  $x = 0.5$
- (ii) Use one application of Newton's method to find a closer approximation to the solution  $x = 0.4$ , correct to three decimal places

**4**

**Question 5** ( Start a new booklet)

**Marks**

- a) In Group A there are 5 men and 3 women. In Group B there are 4 men and 6 women. **4**
- (i) If one person is chosen at random from each group what is the probability that the two people chosen are of opposite sexes?
- (ii) If a group and then one person from that group is chosen at random what is the probability that the person chosen is a man?
- 
- b) Using  $\tan \frac{\theta}{2} = t$  show that  $\frac{1 - \cos \theta}{\sin \theta} = t$  **3**
- 
- c) A particle moves in a straight line such that its acceleration,  $a$ , is given by  $a = 3x^2$ , where  $x$  is displacement,  $v$  is velocity and  $t$  is time. Given that  $v = -\sqrt{2}$  and  $x = 1$  when  $t = 0$  find  $x$  as a function of time,  $t$  **5**

**Question 6** (Start a new booklet)

**Marks**

a) Consider the function  $f(x) = 3 \sin^{-1}\left(\frac{x}{2}\right)$  **5**

- (i) State the domain and range of this function
- (ii) Sketch the graph of  $y = f(x)$
- (iii) Find the slope of the graph at  $x = 0$

b) The displacement,  $x$  metres, of a particle, at  $t$  seconds is given by: **7**

$$x = 5 \cos(4\pi t)$$

- (i) Show that the acceleration of the particle can be expressed in the form:


$$\ddot{x} = -n^2 x$$

- (ii) State the period,  $P$ , of the motion
- (iii) Determine the maximum velocity of the particle
- (iv) Determine the maximum acceleration of the particle
- (v) Express  $v^2$  in terms of  $x$ , where  $v$  is the velocity of the particle

**QUESTION 7 IS ON THE NEXT PAGE**

- a) A vertical building stands with its base  $O$  on horizontal ground.  $A$  and  $B$  are two points on the building vertically above each other such that  $A$  is  $4h$  metres above  $O$  and  $B$  is  $h$  metres above  $O$ . A particle is projected horizontally with speed  $U$  metres per second from  $A$  and 10 seconds later a second particle is projected horizontally with speed  $V$  metres per second from  $B$ . The two particles hit the ground at the same point and at the same time.

**7**



- (i) Show that the horizontal and vertical displacements of particles  $A$  and  $B$ ,  $t$  seconds after the first particle is projected, are given by  $x_A = Ut$ ,  $y_A = 4h - \frac{1}{2}gt^2$  and  $x_B = V(t-10)$ ,  $y_B = h - \frac{1}{2}g(t-10)^2$ , respectively
- (ii) Find the time of flight of each particle
- (iii) Prove that  $V = 2U$
- b) The point  $P(2t, t^2)$  is on the parabola with equation  $x^2 = 4y$ , having its focus at  $F$ . The point  $M$  divides the interval  $FP$  externally in the ratio 3:1.
- (i) Show that the co-ordinates of  $M$  are  $x = 3t$  and  $y = \frac{1}{2}(3t^2 - 1)$
- (ii) Hence prove that the locus of  $M$  is also a parabola and determine the focal length of the locus of  $M$

**5**



## STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

NOTE :  $\ln x = \log_e x, \quad x > 0$