


Mr Antonio
Mrs Collett
Mrs Kerr
Ms Lau
Mrs Soutar

Name:

Teacher:



2011
TRIAL HIGHER SCHOOL CERTIFICATE
EXAMINATION

Mathematics Extension 1

General Instructions

- Reading time – 5 minutes
- Working time – 2 hours
- Write using black or blue pen
- Board-approved calculators may be used
- A table of standard integrals is provided
- All necessary working should be shown in every question
- Start each question in a new booklet
- Marks may be deducted for careless or untidy work

Total Marks – 84

- Attempt Questions 1-7
- All questions are of equal value

Mark	/84
Rank	/
Highest Mark	/84

Blank Page

Question 1. (12 Marks) Use a **SEPARATE** Writing Booklet.

Marks

- (a) Find $\int \frac{2}{\sqrt{36-x^2}} dx$. **1**
- (b) Differentiate $\tan^{-1}(\ln x)$ with respect to x . **1**
- (c) Find the coordinates of the point P that divides the interval joining A (-3, 8) and B (7, -3) internally in the ratio 2 : 3. **2**
- (d) Solve $\frac{4}{x-2} \leq 2$. **3**
- (e) The acute angle between the lines $2x - y = 4$ and $y = mx + 3$ is 45° .
Find the two possible values of m . **2**
- (f) Use the substitution $u = x - 3$ to evaluate $\int_4^5 \frac{x}{\sqrt{x-3}} dx$. **3**

Question 2 (12 Marks) Use a **SEPARATE** Writing Booklet.

Marks

- (a) Let $f(x) = 3 \sin^{-1} 2x$. **2**

Sketch the graph of $y = f(x)$, clearly indicating the endpoints for the domain and the range.

- (b) (i) Differentiate $x \cos^2 x$ with respect to x . **2**

- (ii) Hence, or otherwise, find $\int x \sin 2x \, dx$. **2**

- (c) The polynomial $P(x) = x^3 + ax^2 - 2x + b$ has $(x + 1)$ as a factor. $P(x)$ has a remainder of 4 when divided by $(x - 3)$. **3**

Find the values of a and b .

- (d) The function $f(x) = x - e^{-2x}$ has one root between $x = 0$ and $x = 1$. **3**
Use one application of Newton's method, starting at $x = 0.3$, to find another approximation for this root.

Write your answer correct to 2 decimal places.

Question 3 (12 Marks) Use a SEPARATE Writing Booklet.


Marks

(a) Evaluate $\lim_{x \rightarrow \infty} \frac{3x^2 + 1}{2 - 7x - 5x^2}$. 2

(b) (i) Express $\sqrt{3} \cos \theta - \sin \theta$ in the form $r \cos(\theta + \alpha)$, where $r > 0$, and $0 < \alpha < \frac{\pi}{2}$, giving r and α as exact values. 2

(ii) Solve $\sqrt{3} \cos \theta - \sin \theta = -1$, for $0 \leq \theta \leq 2\pi$, giving your answers as exact values. 2


(c) In the diagram below, AD is perpendicular to BC.
CD = 4, BD = 2 and AD = 3. $\angle CAD = \alpha$ and $\angle BAD = \beta$. 3



Find the exact value of $\sin(\alpha - \beta)$.

- (d) WZ and XY are tangents to the circles WXY and WYZ respectively. The circles share two common points, W and Y. Copy or trace the diagram into your Writing Booklet.

3



Prove that $WX \parallel YZ$.

Question 4 (12 Marks) Use a **SEPARATE** Writing Booklet.

Marks

(a) Find the exact value of $\int_0^{\ln 3} \frac{e^x}{e^x + 9} dx$. 2

(b) Find the constant term in the expansion $\left(2x - \frac{1}{x}\right)^6$. 2

(c) Prove by induction that $4p + 3p^2 + 2p^3$ is divisible by 3 for $p = 1, 2, 3, \dots$ 3

(d) The temperature ($T^\circ\text{C}$) of steel, after it has been removed from a hot furnace, after t minutes, satisfies the differential equation:

$$\frac{dT}{dt} = k(T - 22) \quad \text{where } k \text{ is a constant.}$$

Initially, the temperature (T) of the steel is 100°C and when $t=15$ minutes, $T=70^\circ\text{C}$.

(i) Use this information to find the exact values of A and k . 2

(ii) Hence find the value of t when $T = 40^\circ\text{C}$ to the nearest minute. 1

(e) Show that: 2

$$\sqrt{\frac{1 + \sin 2\theta}{1 - \sin 2\theta}} = \frac{1 + \tan \theta}{1 - \tan \theta}$$

Question 5 (12 marks) Use a **SEPARATE** Writing Booklet.

Marks

(a) The speed V cm/s of a particle moving along the x – axis is given by

$$V^2 = 27 + 18x - 9x^2 \text{ where } x \text{ is in cm.}$$

(i) Prove that the motion is Simple Harmonic. Find the period and amplitude of the motion. 3

(ii) Find the acceleration of the particle when it is 1cm away from the centre of motion. 1

(b) A stone is projected upwards from the edge of a cliff with a speed of 30m/s. It hits an object 120 m horizontally from the edge and 35 m vertically below it.

Assume that t seconds after the release, the position of the stone is given by

$$x = 30t \cos \alpha \text{ and } y = -5t^2 + 30t \sin \alpha .$$

(i) Find α , the angle of projection, to the nearest minute. 3

(ii) Find the time taken for the stone to hit the object. 1


(c) (i) Sketch the curve $y = x + \frac{4}{x}$ showing clearly all stationary points and asymptotes. 3

(ii) Hence, find the values of k such that $x + \frac{4}{x} = k$ has no real roots. 1

Question 6 (12 marks) Use a SEPARATE Writing Booklet.

Marks

(a)



In the above diagram, the points A, B and O are in the same horizontal plane. A and B are 50 m apart and $\angle AOB = 60^\circ$. OT is a vertical tower of height h metres.

The angles of elevation of T from A and B respectively are 45° and α° (α is acute).

- (i) Explain why $AO = h$. 1
- (ii) Prove $h^2 \cot^2 \alpha - h^2 \cot \alpha + h^2 = 50^2$. 2
- (iii) Given that the tower is 30m high, find the angle α correct to the nearest degree. 3

Question 6 - continued.

Marks

- (b) (i) Verify that $\frac{d}{dx}(x \sin^{-1} x + \sqrt{1-x^2}) = \sin^{-1} x$. 1
- (ii) Hence, using a similar expression, find a primitive of $\cos^{-1} x$. 1
- (iii) The curves $y = \sin^{-1} x$ and $\cos^{-1} x$ intersect at P . 1
The curve $y = \cos^{-1} x$ also intersects the x axis at Q .
Show that P has co-ordinates $\left(\frac{1}{\sqrt{2}}, \frac{\pi}{4}\right)$.
- (iv) Find the area enclosed by the x – axis and the arcs OP and PQ . 3

Question 7(12 marks) Use a **SEPARATE** Writing Booklet.

Marks

- (a) A particle is moving in a straight line with acceleration given by

$$\frac{d^2x}{dt^2} = 9(x-2).$$

where x is the displacement in metres, from the origin O after t seconds.

Initially the particle is 4m to the right of O and it has a velocity of $V = 6\text{m/s}$.

- (i) Show that $V^2 = 9(x-2)^2$. **3**
- (ii) Find an expression for V and hence find x as a function of t . **3**
- (b) PQ is a variable chord of the parabola $x^2 = 4ay$ which subtends a right angle at the vertex.
- (i) If p and q are the parameters corresponding to the points P and Q , prove that $pq = -4$. **2**
- (ii) Show that the equation of the normal at P is $x + py = 2ap + ap^3$. **2**
- (iii) Hence prove that the locus of the point of intersection of the normals at P and Q is the parabola $x^2 = 16a(y - 6a)$. **2**

End of Paper

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, \quad x > 0$

