

kl

TABLE #1 Before Editing

Table 1¹ identifier
Body and Lung Weights of Five Lambs title

| Lamb | Body weight (kg) | Dry, Bloodless Lung Weight (g) |
|------------|---------------------|-----------------------------------|
| 15 | 7.3 | 19.7 |
| 16 | 6.9 | 18.9 |
| 17 | 5.4 | 18.0 |
| 18 | 10.0 | 24.6 |
| 19 | 5.2 | 15.5 |
| Mean ± SEM | 7.0±0.9 | 19.3±1.5 |

Abbreviation: SEM, Standard Error of the Mean

TABLE #1 After Editing

Table 1
Body and Lung Weights of Five Normal One Day Old Mixed Breed Lambs

| Lamb | Body Weight (kg) | Lung Weight (g) |
|------------|------------------|-----------------|
| 15 | 7.3 | 19.7 |
| 16 | 6.9 | 18.9 |
| 17 | 5.4 | 18.0 |
| 18 | 10.0 | 24.6 |
| 19 | 5.2 | 15.5 |
| Mean ± SEM | 7.0±0.9 | 19.3±1.5 |

Abbreviation: SEM, Standard Error of the Mean
"Lung Weight" is dry bloodless.

Table 2 Before Editing

Table 2 Identifier
 Data Related to Lung Fluid Balance in Five Anephric, Anesthetized 1-2-wk old Lambs Before and After Intravenous Furosemide
 8 mg/kg ^{Table}

| Lamb | Weight (kg) | Dry, Bloodless Lung (g) | Time Relative to Infusion | Vascular Pressures | | | Protein Concentrations | | | Plasma Column Header |
|------------|-------------|-------------------------|---------------------------|--------------------|------------|--------------------|------------------------|--------------|-------------------|----------------------|
| | | | | Ppa (Torr) | Pla (Torr) | Lymph Total (g/dl) | %Alb ^a | Total (g/dl) | %Alb ^a | |
| 15 | 7.3 | 19.7 | Before | 17 | 7 | 2.90 | 55 | 3.90 | 49 Data | |
| | | | After | 17 | 7 | 2.83 | 55 | 4.04 | 47 | |
| 16 | 6.9 | 18.9 | Before | 17 | 8 | 2.70 | 48 | 4.01 | 45 | |
| | | | After | 17 | 9 | 2.57 | 53 | 3.96 | 44 | |
| 17 | 5.4 | 18.0 | Before | 21 | 7 | 2.40 | 50 | 4.03 | 47 | |
| | | | After | 21 | 7 | 2.60 | 46 | 4.19 | 43 | |
| 18 | 10.0 | 24.6 | Before | 25 | -1 | 2.14 | 55 | 3.78 | 47 | |
| | | | After | 27 | 1 | 2.26 | 54 | 3.92 | 48 | |
| 19 | 5.2 | 15.5 | Before | 21 | 3 | 1.95 | 64 | 3.78 | 52 | |
| | | | After | 21 | 3 | 1.91 | 65 | 3.76 | 52 | |
| Mean ± SEM | 7.0 ± 0.9 | 19.3 ± 1.5 | Before | 20 ± 1 | 5 ± 2 | 2.42 ± 0.17 | 54 ± 3 | 3.90 ± 0.05 | 48 ± 1 | |
| | | | After | 21 ± 2 | 5 ± 1 | 2.43 ± 0.16 | 55 ± 3 | 3.97 ± 0.07 | 47 ± 2 | |
| <i>p</i> | | | | NS | NS | NS | NS | NS | NS | |

^a Percent of total protein which is albumin
 Abbreviations: SEM, Standard Error of the Mean; Ppa, mean pulmonary arterial pressure; Pla, mean left atrial pressure.
 Table #2 After Editing

Table 2 After Editing

Table 2

Comparison of Vascular Pressure and Protein Concentrations Including Percent of Albumin in Lymph and Plasma in Five Anephric 1-2 Week Old Lambs Before and After Intravenous Furosemide Application

| Lamb | Weight Body (kg) | Lung (g) | Time | Vascular Pressures | | Protein Concentrations | | | |
|------------|------------------------|----------|--------|--------------------|---------------|------------------------|-------------------|-------------------|------|
| | | | | Ppa (Torr) | Pla (Torr) | Total | Lymph | Plasma | |
| | | | | | | Total | %Alb ^a | %Alb ^a | |
| 15 | 7.3 | 19.7 | Before | 17 | 7 | 2.90 | 55 | 3.90 | 49 |
| | | | After | 17 | 7 | 2.83 | 55 | 4.04 | 47 |
| 16 | 6.9 | 18.9 | Before | 17 | 8 | 2.70 | 48 | 4.01 | 45 |
| | | | After | 17 | 9 | 2.57 | 53 | 3.96 | 44 |
| 17 | 5.4 | 18.0 | Before | 21 | 7 | 2.40 | 50 | 4.03 | 47 |
| | | | After | 21 | 7 | 2.60 | 46 | 4.19 | 43 |
| 18 | 10.0 | 24.6 | Before | 25 | -1 | 2.14 | 55 | 3.78 | 47 |
| | | | After | 27 | 1 | 2.26 | 54 | 3.92 | 48 |
| 19 | 5.2 | 15.5 | Before | 21 | 3 | 1.95 | 64 | 3.78 | 52 |
| | | | After | 21 | 3 | 1.91 | 65 | 3.76 | 52 |
| Mean ± SEM | 7.0±0.9 | 19.3±1.5 | Before | 20±1 | 5±2 | 2.42±0.17 | 4±3 | 3.90±0.05 | 48±1 |
| | | | After | 21±2 | 5±1 | 2.43±0.16 | 5±3 | 3.97±0.07 | 47±2 |
| <i>p</i> | | | | NS | NS | NS | NS | NS | NS |

^a Percent of total protein which is albumin
 Abbreviations: SEM, Standard Error of the Mean; Ppa, mean pulmonary arterial pressure; Pla, mean left atrial pressure,
 Time is Relative to Infusion, Protein Concentration Total = g/dl.

Table A
Summary of November 1999 Standard Curves and Peak Ratios For Test Drug IERJCMKIRBCM

| Batch | Nov. Run Day | 0.10 | 0.20 | 1.00 | 2.00 | 4.00 | 5.00 | 7.00 |
|-------|--------------|-----------|-------|-----------|-----------|-------|-----------|-----------|
| 1 | 12 | 0.017 R08 | 0.052 | 0.256 | 0.523 | 1.058 | 2.169 R06 | 2.169 R01 |
| 2 | 13 | 0.036 R08 | 0.072 | 0.358 | 0.221 R03 | 0.924 | 1.039 | 1.466 R08 |
| 3 | 15 | 0.019 R08 | 0.038 | 0.221 R03 | | 0.924 | 1.039 | 1.466 R08 |
| 4 | 16 | 0.021 R08 | 0.044 | 0.273 | 0.499 | 1.012 | 1.389 R03 | 1.937 |
| 5 | 17 | 0.023 R03 | 0.053 | 0.250 | 0.484 | 1.069 | 1.240 | 1.569 R08 |
| 6 | 18 | 0.023 R08 | 0.041 | 0.212 | R03 | 0.790 | 1.089 | 1.425 R08 |
| 7 | 19 | 0.021 R08 | 0.041 | 0.221 | 0.455 | 0.929 | 1.140 | 1.548 R08 |
| 8 | 22 | R03 | 0.046 | 0.230 | 0.479 | 0.922 | 1.153 | 1.686 R08 |
| 9 | 23 | 0.026 R08 | 0.042 | 0.232 | 0.409 | 0.843 | 1.141 | R01 |
| 201 | 18 | 0.022 R08 | 0.052 | 0.227 | 0.445 | 0.914 | 1.123 | 1.594 R08 |

Note.

- R01 - Sample or standard lost during the extraction procedure.
- R03 - Internal standard did not meet acceptance criteria for this analyte or this study. Unknowns must be rejected unless otherwise specified in the method report or Study Overview. Standards are knowns and may be rejected.
- R06 - Standard is an outlier: the value is greater than $\pm 15\%$ away from the nominal values, or greater than $\pm 20\%$ away from the nominal value at the LLOQ.
- R08 - Standard not needed to cover study sample range.

Recommendations applied to this edited table:

1. Shorten dates
2. Clarify title
3. Add rule lines
4. Align columns
5. Insure that the data is correct. Some of it was questionable because of alignment