

**Question One (Start a new page)****Marks**

- a) Consider the complex numbers  $z_1 = \sqrt{2}(1 + i\sqrt{3})$  and  $z_2 = 2\sqrt{6}(1 + i)$ .
- i. Express  $z = \frac{z_1}{z_2}$  in the form of  $x + iy$ , where  $x$  and  $y$  are real. **2**
- ii. Write  $z_1$ ,  $z_2$  and  $z$  in modulus/ argument form. **5**
- iii. Hence find the exact value of  $\cos \frac{\pi}{12}$ . **1**
- b) Sketch on separate Argand diagrams the regions where
- i.  $\operatorname{Re}(z + iz) \geq 2$  **2**
- ii.  $1 \leq |z - 1 - i| \leq 3$  where  $z = x + iy$ . **2**
- c) By applying De Moivre's Theorem and by also expanding  $(\cos \theta + i \sin \theta)^5$ , express  $\sin 5\theta$  as a polynomial in  $\sin \theta$ . **3**

**Question Two (Start a new page)**

- a) Given real positive numbers  $a$ ,  $b$  and  $c$  such that  $a > b > c$ .
- i. Prove that  $(a + b) > 2\sqrt{ab}$ . **1**
- ii. Show that  $b^2 - a^2 < 2(b - a)\sqrt{ab}$ . **1**
- iii. Deduce that  $(b - a)\sqrt{a} + (c - b)\sqrt{c} > \frac{c^2 - a^2}{2\sqrt{b}}$ . **2**
- b) i. Sketch the graph of  $f(x) = 1 - \frac{9}{x^2} + \frac{18}{x^4}$ , showing all stationary points and other essential features. **5**
- ii. Hence find the set of values of the real numbers  $k$  such that the equations  $f(x) = k$  has four distinct real roots. **1**
- c) An object of mass  $m$  kg is travelling around a circular banked track of radius  $r$  metres and angle of banking  $\theta$ . The mass is travelling at  $v \text{ ms}^{-1}$ . The forces acting on the object are the gravitational force  $mg$  newtons, a sideways friction force  $F$  newtons (acting down the road as shown) and a normal reaction  $N$  newtons to the road.




Diagram  
not to scale

- i. By resolving forces vertically and horizontally, derive expressions for  $N$  and  $F$ . 4
- ii. Given the radius of the curve is 1 km and  $\tan \theta = \frac{1}{100}$ , find the velocity which will ensure no sideways friction. (Take  $g = 10\text{ms}^{-1}$ ) 1

### Question Three (Start a new page)

- a) Evaluate  $\int_0^4 \frac{dx}{3 + \sqrt{x}}$ . 3
- b) Let  $\alpha, \beta$  and  $\delta$  be the roots of  $x^3 - x^2 + 2x - 1 = 0$ .
  - i. Find the value of  $\alpha + \beta + \delta$ . 1
  - ii. Hence, or otherwise, find the cubic equation with roots :  $-(\alpha + \beta), -(\beta + \delta)$  and  $-(\alpha + \delta)$ . 2
- c) Consider the ellipse  $E$  with equation  $\frac{x^2}{16} + \frac{y^2}{9} = 1$  and its auxillary circle  $C$  with equation  $\frac{x^2}{16} + \frac{y^2}{16} = 1$ .

A straight line  $l$  parallel to the  $y$  axis, intersects the  $x$  axis at  $N$  and the curves  $E$  and  $C$  at the points  $P$  and  $Q$  respectively.

Given that  $P$  and  $Q$  are both in the first quadrant and the coordinates of  $P$  on  $E$  are  $(4 \cos \theta, 3 \sin \theta)$ .

- i. Sketch the curves  $E$  and  $C$  showing the above information. 1
- ii. Write down the coordinates of  $N$  and  $Q$  in terms of  $\theta$ . 2
- iii. Derive the equation of the tangent to the curve  $E$  at the point  $P$ . 2
- iv. Write down the equation of the tangent to the curve  $C$  at the point  $Q$ . 1
- v. The tangents at  $P$  and  $Q$  intersect at a point  $R$ . Show that  $R$  lies on the  $x$  axis. 2
- vi. Prove that  $ON \cdot OR$  is independent of the positions  $P$  and  $Q$ . 1

**Question Four (Start a new page)**

**Marks**


a) Using the Table of Standard Integrals, find  $\int \frac{1}{\sqrt{x^2 - 4x + 5}} dx$ . 2

b) i. Find the constants  $A$  and  $B$  such that 2

$$\frac{1}{\cos x} = \frac{A \cos x}{1 - \sin x} + \frac{B \cos x}{1 + \sin x}.$$

ii. Hence find the exact value of  $\int_0^{\frac{\pi}{6}} \sec x dx$ . 2

c) *Diagram not to scale*



In the above diagram,  $C$  is a circle with exterior point  $T$ . Tangents from  $T$  are drawn to meet  $C$  at the points  $A$  and  $E$ . The point  $O$  is the centre of  $C$ . The line  $BT$  passes through  $O$ . The line  $AD$  passes through  $B$ . The line  $OF$  passes through  $E$ .  $AT$  is parallel to  $DF$ .

i. Trace or copy the diagram onto your answer book and prove  $\triangle OET \cong \triangle OAT$ . 2

ii. Considering  $\triangle OET$  and  $\triangle DEF$ , show that  $DE = \frac{DF(ET^2 - OE^2)}{OT^2}$  by using double angle formula. 4

iii. Use the sine rule to show that  $\frac{AB}{BD} = \frac{AT}{DT}$ . 3

**Question Five (Start a new page)****Marks**

- a) i. Prove  $\int_0^a f(x)dx = \int_0^a f(a-x)dx$ . **2**
- ii. Hence evaluate  $\int_0^1 x(1-x)^n dx$ , . **3**
- b) Let  $w = \cos \frac{\pi}{5} + i \sin \frac{\pi}{5}$ .
- i Find all the complex roots of the equation  $z^{10} - 1 = 0$  and write them down in terms of  $w$  and  $k$ , while  $k$  is a positive integer. **2**
- ii. Prove that  $1 + w + w^2 + w^3 + \dots + w^9 = 0$ . **1**
- iii. The quadratic equation  $x^2 + bx + c = 0$ , where  $b$  and  $c$  are real, has the root  $w + w^4$ . Find the other root in terms of  $w$ . **2**
- iv. Find  $b$  and express  $c$  in terms of  $\sin \frac{\pi}{5}$ . **5**

**Question Six (Start a new page)**

- a) Find  $\int \frac{1}{e^x + e^{-x}} dx$ . **3**
- b) Given that  $I_n = \int_1^e (\ln y)^n dy$ ,  $n = 0, 1, 2, 3, \dots$
- i. Prove that  $I_n = e - nI_{n-1}$  **3**
- ii. Hence evaluate  $\int_1^e (\ln y)^2 dy$ . **2**
- c) The depth of water at the entrance to a harbour can be modeled using the equation  $x = b + a \cos nt$  where  $x$  metres is the depth of water and  $t$  is time measured in hours. For a certain harbour, the first low tide for the day is at 5am and the water depth is 20m. The next high tide is  $6\frac{1}{2}$  hours later and the corresponding depth is 28m.
- i. Taking the first low tide for the day as the origin for measuring the time, write down the values of  $a$ ,  $b$  and  $n$ . **2**
- ii. Find the depth of water at 9am. (correct to 3 significant figures) **2**
- iii. Find all the times after mid-night and before mid-day when water depth is 23 m. (correct to nearest minute) **2**
- iv. Find the greatest rate at which the tide is rising. **1**


**Question Seven (Start a new page)****Marks**

- a) The circle  $x^2 + y^2 = 9$  is rotated about the line  $x = 8$  to form a torus. Using the method of cylindrical shells, find the volume of the torus. **4**
- b) i.  $xy = c^2$  is the result of rotating  $x^2 - y^2 = a^2$  anticlockwise through an angle of  $45^\circ$ . Write down the relationship between  $a^2$  and  $c^2$ . **1**
- ii.  $P(x_1, y_1)$  is the point of intersection of the hyperbolas  $xy = c^2$  and  $x^2 - y^2 = a^2$  in the first quadrant. Prove that the tangent to  $xy = c^2$  at the point  $P$  is  $xy_1 + yx_1 = 2c^2$ . **2**
- iii. Write down the equation of the tangent to  $x^2 - y^2 = a^2$  at the point  $P$ . **1**
- iv. The tangent to the hyperbola  $x^2 - y^2 = a^2$  at  $P$  meets its asymptotes at  $A$  and  $C$  while the tangent to the hyperbola  $xy = c^2$  at  $P$  meets its asymptotes in  $B$  and  $D$ .
- v. Show that the co-ordinates of  $A$  are  $(x_1 + y_1, x_1 + y_1)$ . **2**
- vi. Find the co-ordinates of  $B, C$  and  $D$ . **3**
- vii. Prove that  $ABCD$  is a square. **2**

**Question Eight (Start a new page)**

- a) As shown in the diagram below, a light string of  $2l$  metres long is attached to two points  $A$  and  $B$ . A mass of  $3m$  kg is attached to the middle of the string and a second mass of  $m$  kg in the form of a ring is attached to the end of the string at  $B$ .


The  $3m$  kg mass is rotating in circular motion at  $\omega$  radians per second and the  $m$  kg mass is free to move up or down the smooth vertical rod  $AB$ . The string makes an angle  $\theta$  with the vertical. (Assume the acceleration due to gravity is  $g \text{ ms}^{-2}$ ).



*Diagram not to scale*

- i. Given that  $h$  is the distance between  $A$  and the centre of the circular motion, find an expression for  $h$  in terms of  $g$  and  $\omega$ . 4
- ii. If the  $3m$  kg mass is replaced by a mass of  $m$  kg mass and the  $m$  kg ring is replaced by a ring of  $3m$  kg, the speed of the rotating mass is doubled to  $2\omega$  radians per second. Determine if  $h$  is increased or decreased and give reasons. (note that  $\omega > 1$ ) 3

- b) Mr Dud's crystal ball rests on a solid stand which is in the shape of a square based frustum as shown.



*Diagram not to scale*

The stand is constructed so that the crystal ball of radius  $R$  fits snugly inside and just touches the centre of the square base. The side  $BC$  of the base slopes so that if extended it would pass through the top-most point of the ball at  $A$  and makes an angle  $\theta$  with the vertical  $AD$ . Take  $O$  as the centre of the circle and let the distance  $OL$  be  $x$  units.

- i. Explain why  $LQ = \sqrt{R^2 - x^2}$  and  $LM = (R + x) \tan \theta$ . 2
- ii. Consider a slice  $KLM$  of thickness  $\Delta x$  as shown perpendicular to  $AD$ . 2
- Show that it has a volume  $\Delta V \approx \left\{ 4 \tan^2 \theta (R + x)^2 - \pi (R^2 - x^2) \right\} \Delta x$ .
- iii. Find the volume of such a solid when the angle  $\theta = \frac{\pi}{6}$ . 4

**END**

