

St George Girls' High School

Trial Higher School Certificate Examination

2003



Mathematics

Extension 2

General Instructions

- Reading time – 5 minutes
- Working time – 3 hours
- Write using blue or black pen
- All questions may be attempted.
- Begin each question on a new page
- All necessary working must be shown.
- Marks may be deducted for careless or poorly presented work.
- Board-approved calculators may be used.
- A list of standard integrals is included at the end of this paper.
- The mark allocated for each question is listed at the side of the question.

Students are advised that this is a Trial Examination only and does not necessarily reflect the content or format of the Higher School Certificate Examination.

Question 1 – (15 marks) – Start a new page

Marks

- a) Show that $y = xe^{-x}$ has a stationary point at the point $\left(1, \frac{1}{e}\right)$. 2

- b) On separate diagrams sketch the following curves. Indicate clearly any turning points, asymptotes and intercepts with the coordinate axes. 9

(i) $y = xe^{-x}$

(ii) $y = x^2 e^{-2x}$

(iii) $y = \frac{1}{x^2 e^{-2x}}$

(iv) $y = \log_e(xe^{-x})$

(v) $y = e^{xe^{-x}}$

- c) Solve for x :

$$\frac{3-x}{\sqrt{x-1}} \geq \frac{\sqrt{x-1}}{3-x}$$

4

Question 2 – (15 marks) – Start a new page

a) Evaluate the following definite integrals:

(i) $\int_0^1 \frac{2x}{1+2x} dx$ 3

(ii) $\int_2^3 \frac{x+1}{\sqrt{x^2+2x+5}} dx$ 3

(iii) $\int_2^4 \frac{dx}{x\sqrt{x-1}}$ 3

b) The cubic equation $x^3 - 2x + 4 = 0$ has roots α, β, γ .

(i) Prove that the cubic equation with roots $\alpha^2, \beta^2, \gamma^2$ is $x^3 - 4x^2 + 4x - 16 = 0$. 3

(ii) Factorise $x^3 - 4x^2 + 4x - 16$ into linear factors and hence prove that only one of α, β, γ is real. 3

Question 3 – (15 marks) – Start a new page

Marks

a) Let $P(z) = z^7 - 1$.

(i) Find all the complex roots of $P(z) = 0$. (Call them z_0, z_1, \dots, z_6 leaving all answers in terms of π). 3

(ii) Plot the points representing z_0, z_1, \dots, z_6 on the Argand Diagram.

(iii) Factorise $P(z)$ over the complex numbers. 1

(iv) Factorise $P(z)$ over the real numbers (leave answers in terms of π). 2

(v) Show that $\cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{6\pi}{7} = -\frac{1}{2}$ 2

b) The area bounded by the curve $y = x(2 - x)$ and the x axis is rotated about the y axis. 5

(i) By considering cylindrical shells with generators parallel to the y axis, show that the volume V units³ of the solid so generated is given by $V = \int_0^2 2\pi xy \, dx$. (shaded circle)

(ii) Hence, determine the volume of this solid. (shaded circle)

Question 4 – (15 marks) – Start a new page

Marks

a) (i) Prove that $\int_0^a f(x) dx = \int_0^a f(a-x) dx$

2

- (ii) With reference to an appropriate diagram, show geometrically why the above result is true.

2

(iii) Evaluate $\int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$

3

b) If $I_n = \int_0^1 x^n e^{-x} dx \quad (n \geq 0)$

(i) Prove that $I_n = n I_{n-1} - \frac{1}{e} \quad (n \geq 1)$

2

(ii) Hence, evaluate $\int_0^1 x^3 e^{-x} dx$

2

c) Using the substitution $t = \tan \frac{\theta}{2}$, or otherwise, calculate $\int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \cot \theta \tan \frac{\theta}{2} d\theta$

4

Question 5 – (15 marks) – Start a new page

Marks

- a) Find all pairs of real numbers x and y that satisfy $(x + iy)^2 = -12 + 16i$ 3

- b) Consider the polynomial equation

$$z^3 + (6+i)z^2 + (17+8i)z + 33i + 30 = 0$$

- (i) This equation has a root $\alpha = pi$ where p is real. Find the value of p . 3

- (ii) If the other roots are β and γ , use the relationships for the sum and product of the roots (or otherwise) to find β and γ .
[Hint: Use your answer from part (a)]. 3

- (iii) Show that the points representing α, β, γ on the Argand diagram are the vertices of a right-angled triangle. 1

- c) The complex number z and its conjugate \bar{z} satisfy

$$(z - \bar{z})^2 + 8(z + \bar{z}) = 16$$

5


- (i) Prove that the point which represents z on the Argand diagram lies on a parabola.

- (ii) Sketch the locus of z and hence show that $-\frac{\pi}{4} \leq \arg z \leq \frac{\pi}{4}$

Question 6 – (15 marks) – Start a new page

Marks

a)



The point $P(x_1, y_1)$ lies on the ellipse $\frac{x^2}{9} + \frac{y^2}{5} = 1$ ($x_1 \neq 0$)

The tangent to the ellipse at P cuts the x axis at T .

(i) Find the eccentricity, e , of the ellipse.

1

(ii) Find the coordinates of the foci, S and S'

1

(iii) Find the equations of the directrices.

1

(iv) Show that the equation of the tangent at P is $\frac{x \cdot x_1}{9} + \frac{y \cdot y_1}{5} = 1$

2

(v) Hence write down the coordinates of T .


1

(vi) Using the focus-directrix definition of the ellipse, or otherwise, show that

4

$$\frac{PS}{PS'} = \frac{TS}{TS'}$$

b)



NOT TO SCALE

5

$ABCD$ and $AEFC$ are parallelograms. $AE = 5\text{cm}$ and $EB = 3\text{cm}$.

(i) Find, giving brief reasons, the length of AG .

(ii) Find the ratio of the area of $\triangle CDH$ to the area of the parallelogram $ABCD$.

Question 7 – (15 marks) – Start a new page

Marks

- a) The acceleration of a particle which is moving along the x axis is given by

$$\frac{d^2x}{dt^2} = 2x^3 - 10x$$

- (i) If the particle starts at the origin with velocity u show that its velocity v is given by $v^2 - u^2 = x^4 - 10x^2$

2

- (ii) (α) If $u = 3$ show that the particle oscillates within the interval $-1 \leq x \leq 1$.

4


- (β) Is this an example of simple harmonic motion? Give a clear reason for your answer.

1

- (iii) If $u = 6$ carefully describe the motion of the particle.

3

b)



$$x = 3 - y$$

The area enclosed by the curve $y = 4 - x^2$, the line $y = 3x$ and the x axis (for $x \geq 0$) is rotated about the line $x = 3$.

Calculate the volume of the solid generated.

5

Question 8 – (15 marks) – Start a new page

- a) (i) x, y and z are positive integers. Show that if x is a factor of both y and z then x is a factor of $z - y$. 1

(ii) Show that $2^{2^{k+1}} = (2^{2^k})^2$ 1


(iii) $F_n = 2^{2^n} + 1$ defines a set of positive integers called Fermat numbers for $n = 0, 1, 2, \dots$ i.e. $F_0 = 2^{2^0} + 1 = 2^1 + 1 = 3$ 5

Use mathematical induction to show that

$$F_n = F_0 F_1 F_2 \dots F_{n-1} + 2 \text{ for } n \geq 1$$

- (iv) Hence, or otherwise, show that the highest common factor of any two Fermat numbers is 1. 3

[Hint: Let k be a common factor of F_m and F_n , where $m < n$, and use (i) and (iii) to show that $k = 1$]



The graph of $y = f(t)$ $0 \leq t \leq 9$ is shown. If $F(x) = \int_0^x f(t) dt$ find:

- (i) the values of x for which $F(x) = 0$
 (ii) the coordinates of any stationary points.



STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE : $\ln x = \log_e x, \quad x > 0$








Question 1

(a) $y = xe^{-x}$
 $\frac{dy}{dx} = 1 \cdot e^{-x} + x \cdot (-e^{-x})$
 $= e^{-x}(1-x)$

Stat pts when $\frac{dy}{dx} = 0$

$$e^{-x}(1-x) = 0$$
 $x = 1$
 $y = 1 \cdot e^{-1} = \frac{1}{e}$

$(1, \frac{1}{e})$ is a stat pt



$$(c) \frac{3-x}{\sqrt{x-1}} \geq \frac{\sqrt{x-1}}{3-x} \quad x > 1$$


$$\frac{3-x}{\sqrt{x-1}} \geq \frac{(x-1)^2}{3-x}$$

$$\frac{3-x}{1} - \frac{x-1}{3-x} \geq 0$$

$$\frac{9-6x+x^2-x+1}{3-x} \geq 0 \quad (x(3-x))^2$$

$$(x^2 - 7x + 10)(3-x) \geq 0$$

$$(x-2)(x-5)(3-x) \geq 0$$



$$1 < x \leq 2 \text{ or } 3 < x \leq 5$$

Question 2

$$\begin{aligned}
 (a) (i) & \int_0^1 \frac{2x}{1+2x} dx \\
 &= \int_0^1 \frac{1+2x-1}{1+2x} dx \\
 &= \int_0^1 1 - \frac{1}{1+2x} dx \\
 &= \left[x - \frac{1}{2} \ln(1+2x) \right]_0^1 \\
 &= \left(1 - \frac{1}{2} \ln 3 \right) - \left(0 - \frac{1}{2} \ln 1 \right) \\
 &= 1 - \frac{1}{2} \ln 3
 \end{aligned}$$

$$\begin{aligned}
 (ii) & \int_2^3 \frac{x+1}{\sqrt{x^2+2x+5}} dx \\
 &= \frac{1}{2} \int_2^3 (2x+2)(x^2+2x+5)^{-\frac{1}{2}} dx \\
 &= \frac{1}{2} \left[\frac{(x^2+2x+5)^{\frac{1}{2}}}{\frac{1}{2}} \right]_2^3 \\
 &= \left[\sqrt{x^2+2x+5} \right]_2^3 \\
 &= \sqrt{20} - \sqrt{13}
 \end{aligned}$$

$$\begin{aligned}
 (iii) & \int_2^4 \frac{dx}{x\sqrt{x-1}} \\
 &= \int_1^{\sqrt{3}} \frac{2u du}{(u^2+1) \cdot u} \\
 &= 2 \int_1^{\sqrt{3}} \frac{du}{u^2+1} \\
 &= 2 \left[\tan^{-1} u \right]_1^{\sqrt{3}}
 \end{aligned}$$

$$= 2 (\tan^{-1} \sqrt{3} - \tan^{-1} 1)$$

$$= 2 \left(\frac{\pi}{3} - \frac{\pi}{4} \right)$$

$$= \frac{\pi}{6}$$

(a) $x^3 - 2x + 4 = 0$ has roots α, β, γ .

$$\text{Let } P(x) = x^3 - 2x + 4$$

Egⁿ with roots $\alpha^2, \beta^2, \gamma^2$ is

$$P(\sqrt{x}) = 0$$

$$(\sqrt{x})^3 - 2\sqrt{x} + 4 = 0$$

$$x\sqrt{x} - 2\sqrt{x} + 4 = 0$$

$$(x-2)\sqrt{x} = -4$$

$$(x-2)^2 \cdot x = 16$$

$$x^3 - 4x^2 + 4x = 16$$

$$x^3 - 4x^2 + 4x - 16 = 0$$

$$(iii) x^3 - 4x^2 + 4x - 16$$

$$= x^2(x-4) + 4(x-4)$$

$$= (x-4)(x^2+4)$$

$$= (x-4)(x+2i)(x-2i)$$

Since $x^3 - 4x^2 + 4x - 16 = 0$ has roots $\alpha^2, \beta^2, \gamma^2$ then $(x-\alpha^2)(x-\beta^2)(x-\gamma^2)$

$$= (x-4)(x+2i)(x-2i)$$

Thus, without loss of generality,

$$\alpha^2 = 4 \quad \beta^2 = -2i \quad \gamma^2 = 2i$$

i.e only α can be real, β and γ are non-real complex numbers

Question 3

$$(a) P(z) = z^7 - 1$$

$$(i) z^7 - 1 = 0$$

$$z^7 = 1 \Rightarrow |z| = 1$$

Let $z = \cos \theta + i \sin \theta$

$$(\cos \theta + i \sin \theta)^7 = 1$$

$$\cos 7\theta + i \sin 7\theta = 1$$

$$\cos 7\theta = 1 \quad \sin 7\theta = 0$$

$$7\theta = 2k\pi$$

$$\theta = \frac{2k\pi}{7}$$

$$k=0 \quad z_0 = \cos 0 + i \sin 0 = 1$$

$$k=1 \quad z_1 = \cos \frac{2\pi}{7} + i \sin \frac{2\pi}{7}$$

$$k=2 \quad z_2 = \cos \frac{4\pi}{7} + i \sin \frac{4\pi}{7}$$

$$k=3 \quad z_3 = \cos \frac{6\pi}{7} + i \sin \frac{6\pi}{7}$$

$$k=4 \quad z_4 = \cos \frac{8\pi}{7} + i \sin \frac{8\pi}{7}$$

$$= \cos \frac{6\pi}{7} - i \sin \frac{6\pi}{7}$$


$$k=5 \quad z_5 = \cos \frac{10\pi}{7} + i \sin \frac{10\pi}{7}$$

$$= \cos \frac{4\pi}{7} - i \sin \frac{4\pi}{7}$$

$$k=6 \quad z_6 = \cos \frac{12\pi}{7} + i \sin \frac{12\pi}{7}$$

$$= \cos \frac{2\pi}{7} - i \sin \frac{2\pi}{7}$$

(ii)



$$(iii) P(z) = (z-1)(z-z_1)(z-z_2)(z-z_3)(z-z_4) \times (z-z_5)(z-z_6)$$

where z_1, \dots, z_6 are indicated in (i)

(iv)

$$(z-z_1)(z-z_6) = z^2 - (z_1+z_6)z + z_1 z_6$$

$$= z^2 - 2 \cos \frac{2\pi}{7} z + 1$$

Similarly we find $(z-z_2)(z-z_5)$ and $(z-z_3)(z-z_4)$

$$\therefore P(z) =$$

$$(z-1)(z^2 - 2 \cos \frac{2\pi}{7} z + 1)(z^2 - 2 \cos \frac{4\pi}{7} z + 1) \times (z^2 - 2 \cos \frac{6\pi}{7} z + 1)$$


$$(iv) z_0 + z_1 + \dots + z_6 = -\frac{\text{coeff } z^6}{\text{coeff } z^7} = 0$$

$$\therefore 1 + 2 \cos \frac{2\pi}{7} + 2 \cos \frac{4\pi}{7} + 2 \cos \frac{6\pi}{7} = 0$$


$$2(\cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{6\pi}{7}) = -1$$

$$\cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{6\pi}{7} = -\frac{1}{2}$$

(b)



When the cylindrical shell of radius $r = x$, height $h = y$ and width δx is cut parallel to y -axis and "flattened out" its volume δV is approx that of a rectangular prism as shown:



$$\delta V = 2\pi x y \delta x$$

$$V = \lim_{\delta x \rightarrow 0} \sum_{x=0}^2 2\pi x y \delta x = \int_0^2 2\pi x y dx$$

$$(ii) xy = x \cdot x(2-x) = 2x^2 - x^3$$

$$V = 2\pi \int_0^2 2x^2 - x^3 dx$$


$$= 2\pi \left[\frac{2x^3}{3} - \frac{x^4}{4} \right]_0^2$$

$$= 2\pi \left(\frac{16}{3} - \frac{16}{4} - 0 \right)$$

$$\text{Volume} = \frac{8\pi}{3} \text{ units}^3$$

Question 4

$$\begin{aligned}
 (a) (i) & \int_0^a f(a-x) dx \quad \text{Let } u = a-x \\
 & du = -dx \\
 & dx = -du \\
 & x=0 \quad u=a \\
 & x=a \quad u=0 \\
 & = \int_a^0 f(u) \cdot (-1) du \\
 & = \int_0^a f(u) du \\
 & = \int_0^a f(x) dx
 \end{aligned}$$



If $y = f(x)$ $0 \leq x \leq a$ is reflected in the y axis we get

$$y = f(-x) \quad -a \leq x \leq 0$$

If this is then translated ' a ' units to the right we get

$$y = f(-(x-a)) \quad 0 \leq x \leq a$$

$$= f(a-x) \quad 0 \leq x \leq a$$

From the graph it can be seen that the area under $y = f(x)$ from $x=0$ to $x=a$ ($\int_0^a f(x) dx$) equals the area under $y = f(-x)$ from $x=-a$ to $x=0$ and this is in turn equal to the area under $y = f(a-x)$ from $x=0$ to $x=a$ (ie $\int_0^a f(a-x) dx$)

$$\begin{aligned}
 (iii) & \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \\
 & = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin(\frac{\pi}{2}-x)}}{\sqrt{\sin(\frac{\pi}{2}-x)} + \sqrt{\cos(\frac{\pi}{2}-x)}} dx \quad (\text{by (i)})
 \end{aligned}$$

$$= \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx$$

Now

$$\begin{aligned}
 & \int_0^{\frac{\pi}{2}} \left(\frac{\sqrt{\sin x}}{\sqrt{\cos x} + \sqrt{\sin x}} + \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} \right) dx \\
 & = \int_0^{\frac{\pi}{2}} 1 dx \\
 & = [x]_0^{\frac{\pi}{2}} = \frac{\pi}{2} - 0 = \frac{\pi}{2} \\
 \text{Since } & \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \\
 & = \frac{\pi}{4}
 \end{aligned}$$

$$(i) I_n = \int_0^1 x^n e^{-x} dx \quad n \geq 0$$

$$\begin{aligned}
 & = \int_0^1 \frac{d(e^{-x})}{dx} \cdot x^n dx \\
 & = \left[-e^{-x} \cdot x^n \right]_0^1 - \int_0^1 -e^{-x} \cdot n x^{n-1} dx \\
 & = (-e^{-1} \cdot 1 - 0) + n \int_0^1 x^{n-1} e^{-x} dx \\
 & = n I_{n-1} - \frac{1}{e}
 \end{aligned}$$

$$\begin{aligned}
 (ii) \int_0^1 x^3 e^{-x} dx & = I_3 \\
 & = 3I_2 - \frac{1}{e} \\
 & = 3(2I_1 - \frac{1}{e}) - \frac{1}{e} \\
 & = 6(1 \cdot I_0 - \frac{1}{e}) - \frac{4}{e} \\
 & = 6I_0 - \frac{10}{e}
 \end{aligned}$$

$$\begin{aligned}
 I_0 & = \int_0^1 e^{-x} dx \\
 & = \left[-e^{-x} \right]_0^1 \\
 & = -e^{-1} - -e^0 \\
 & = 1 - \frac{1}{e}
 \end{aligned}$$

$$\begin{aligned}
 \int_0^1 x^3 e^{-x} dx & = 6\left(1 - \frac{1}{e}\right) - \frac{10}{e} \\
 & = 6 - \frac{16}{e}
 \end{aligned}$$

Question 5

$$(c) \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \omega t \theta \tan \frac{\theta}{2} d\theta$$

$$t = \tan \frac{\theta}{2} \quad \theta = 2 \tan^{-1} t \\ d\theta = \frac{2}{1+t^2} dt$$

$$\theta = \frac{\pi}{3} \quad t = \tan \frac{\pi}{6} = \frac{1}{\sqrt{3}} \\ \theta = \frac{\pi}{2} \quad t = \tan \frac{\pi}{4} = 1$$

$$\cot \theta = \frac{1-t^2}{2t}$$

$$\int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \omega t \theta \cdot \tan \frac{\theta}{2} d\theta$$

$$= \int_{\frac{1}{\sqrt{3}}}^1 \frac{1-t^2}{2t} \cdot t \cdot \frac{2}{1+t^2} dt$$

$$= \int_{\frac{1}{\sqrt{3}}}^1 \frac{1-t^2}{1+t^2} dt$$

$$= \int_{\frac{1}{\sqrt{3}}}^1 \frac{2-(1+t^2)}{1+t^2} dt$$

$$= \int_{\frac{1}{\sqrt{3}}}^1 \frac{2}{1+t^2} - 1 dt$$

$$= \left[2\tan^{-1}t - t \right]_{\frac{1}{\sqrt{3}}}^1$$

$$= \left(2\tan^{-1}1 - 1 \right) - \left(2\tan^{-1}\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{3}} \right)$$

$$= \left(2 \times \frac{\pi}{4} - 1 \right) - \left(2 \times \frac{\pi}{6} - \frac{1}{\sqrt{3}} \right)$$

$$= \frac{\pi}{6} - 1 + \frac{1}{\sqrt{3}}$$

$$(a) (x+iy)^2 = -12+16i \quad (x, y \text{ real})$$

$$x^2 - y^2 + 2ixy = -12+16i \\ x^2 - y^2 = -12 \quad \text{--- (1)}$$

$$2xy = 16 \\ y = \frac{8}{x} \quad \text{--- (2)}$$

Subst (2) in (1)

$$x^2 - \frac{64}{x^2} = -12$$

$$x^4 + 12x^2 - 64 = 0$$

$$(x^2 + 16)(x^2 - 4) = 0$$

$$x = 2, -2 \quad (x \text{ real})$$

$$y = 4, -4$$

$$x = 2, y = 4 \quad \text{or} \quad x = -2, y = -4$$

$$(k) z^3 + (6+i)z^2 + (17+8i)z + 33i + 30 = 0$$

(i) $\alpha = \rho i$ is a root

$$(\rho i)^3 + (6+i)(\rho i)^2 + (17+8i)\rho i + 33i + 30 = 0$$

$$-\rho^3 i^3 - 6\rho^2 i^2 - i\rho^2 + 17\rho i - 8\rho + 33i + 30 = 0$$

Equating real and imag. parts

$$-\rho^3 - \rho^2 + 17\rho + 33 = 0 \quad \text{--- (1)}$$

$$-6\rho^2 - 8\rho + 30 = 0 \quad \text{--- (2)}$$

$$\text{From (2)} \quad 3\rho^2 + 4\rho - 15 = 0$$

$$(3\rho - 5)(\rho + 3) = 0$$

$$\rho = \frac{5}{3}, -3$$

Test in (1)

$$\rho \neq \frac{5}{3} \text{ since } 5 \times 33 + 3 \times 1$$

$$\rho = -3 \quad \text{LHS} = -(-3)^3 - (-3)^2 + 17 \times -3 + 33 \\ = 27 - 9 - 51 + 33$$

$$= 0 = \text{RHS}$$

$$\therefore \alpha = -3i$$

$$(ii) \alpha + \beta + \gamma = -(6+i)$$

$$-3i + \beta + \gamma = -6-i$$

$$\beta + \gamma = -6 + 2i \quad \text{--- (1)}$$

$$\alpha \beta \gamma = -(33i + 30)$$

$$-3i \beta \gamma = -3i(11 - 10i)$$

$$\beta \gamma = 11 - 10i \quad \text{--- (2)}$$

$$\text{From } \textcircled{1} \quad \gamma = -6+2i - \beta$$

Subst in \textcircled{2}

$$\beta(-6+2i - \beta) = 11-10i$$

$$(-6+2i)\beta - \beta^2 = 11-10i$$

$$\beta^2 + (6-2i)\beta + 11-10i = 0$$

$$\beta^2 + (6-2i)\beta + (3-i)^2 = -11+10i + (3-i)$$

$$(\beta + (3-i))^2 = -11+10i + 9-6i - 1 \\ = -3+4i$$


$$\beta + (3-i) = \pm \sqrt{-3+4i} \quad (*) \\ = \pm (1+2i)$$

$$\beta = -3+i + 1+2i, -3+i - 1-2i \\ = -2+3i, -4-i$$

$$\star \sqrt{-12+16i} = 2\sqrt{-3+4i}$$

$$\pm (2+4i) = 2\sqrt{-3+4i}$$

$$\therefore \sqrt{-3+4i} = \pm (1+2i)$$



$$\text{Grad } BC = \frac{3-(-1)}{-2-4}$$

$$= \frac{4}{-2} \\ = -2$$

$$\text{Grad } AC = \frac{-1-(-3)}{-4-0}$$

$$= \frac{2}{-4} \\ = -\frac{1}{2}$$

$$\text{Grad } BC \times \text{Grad } AC = 2 \times -\frac{1}{2} = -1$$

$\therefore A, B, C$ vertices of right-angled A

$$(c) (\bar{z} - \bar{\beta})^2 + 8(z + \bar{\beta}) = 16$$

$$\text{Let } z = x + iy$$

$$\therefore \bar{z} = x - iy$$

$$z - \bar{z} = 2iy$$


$$z + \bar{z} = 2x$$

\therefore Eqn becomes

$$(2iy)^2 + 8 \cdot 2x = 16$$

$$-4y^2 + 16x = 16 \\ 16x = 16 + 4y^2 \\ x = 1 + \frac{y^2}{4}$$

which is a parabola.



Maximum and minimum values of $\arg z$ occur when $y = mx$ is a tangent to $x = 1 + \frac{y^2}{4}$

$$y = mx \quad \text{--- (1)}$$

$$x = 1 + \frac{y^2}{4} \quad \text{--- (2)}$$

Subst (1) in (2)

$$x = 1 + \frac{(mx)^2}{4}$$

$$4x = 4 + m^2x^2$$

$$m^2x^2 - 4x + 4 = 0$$

Line is a tangent when quadratic has equal roots i.e. $\Delta = 0$

$$(-4)^2 - 4 \cdot m^2 \cdot 4 = 0$$

$$16 - 16m^2 = 0$$

$$m^2 = 1$$

$$m = \pm 1$$

$$\tan \theta = \pm 1$$

$$\theta = \pm \frac{\pi}{4}$$

$$\therefore -\frac{\pi}{4} \leq \arg z \leq \frac{\pi}{4}$$

Question 6

$$(a) \frac{x^2}{9} + \frac{y^2}{5} = 1$$

$$a^2 = 9 \quad b^2 = 5 \\ a = 3 \quad b = \sqrt{5}$$

$$(i) b^2 = a^2(1-e^2)$$

$$5 = 9(1-e^2)$$

$$e^2 = \frac{4}{9}$$

$$e = \frac{2}{3} \quad (e > 0)$$

$$(ii) ae = 3 \times \frac{2}{3} = 2$$

$$S(2,0) \quad S'(-2,0)$$

$$(iii) \frac{a}{e} = \frac{3}{\frac{2}{3}} = \frac{9}{2}$$

$$\text{Directrices: } x = \frac{9}{2}, \quad x = -\frac{9}{2}$$

$$(iv) \frac{x^2}{9} + \frac{y^2}{5} = 1$$

$$\frac{2x}{9} + \frac{2y}{5} \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = -\frac{2x}{9} \times \frac{5}{2y} \\ = -\frac{5x}{9y}$$

$$\text{At } P(x_1, y_1) \quad \frac{dy}{dx} = -\frac{5x_1}{9y_1}$$

Eqn of tangent at P is

$$y - y_1 = -\frac{5x_1}{9y_1}(x - x_1)$$

$$\frac{yy_1}{5} - \frac{y_1^2}{5} = -\frac{xx_1}{9} + \frac{x_1^2}{9}$$

$$\frac{xx_1}{9} + \frac{yy_1}{5} = \frac{x_1^2}{9} + \frac{y_1^2}{5} \\ = 1 \quad (\text{since } P(x_1, y_1) \text{ lies on ellipse})$$

$$\frac{xx_1}{9} + \frac{yy_1}{5} = 1$$

(v) When $y = 0$


$$\frac{xx_1}{9} = 1$$

$$x = \frac{9}{x_1}$$

$$\therefore T \text{ has coords } \left(\frac{9}{x_1}, 0\right)$$

$$M\left(\frac{9}{2}, y_1\right)$$

(vi)




Let $M\left(\frac{9}{2}, y_1\right)$ and $M'\left(-\frac{9}{2}, y_1\right)$ be the feet of the perpendiculars from P to the directrices

$$\frac{PS}{PM} = \frac{PS'}{PM'} \quad (\text{focus-directrix definition})$$

$$\therefore \frac{PS}{PS'} = \frac{PM}{PM'} = \frac{\frac{9}{2} - x_1}{\frac{9}{2} + x_1} \\ = \frac{9 - 2x_1}{9 + 2x_1}$$

$$\text{Now } \frac{TS}{TS'} = \frac{\frac{9}{2} - 2}{\frac{9}{2} + 2} \\ = \frac{9 - 2x_1}{9 + 2x_1} \\ = \frac{PS}{PS'}$$



Let $CB = x$

In $\triangle BAC$

$$\frac{BH}{HC} = \frac{BE}{EA} \quad (\text{line parallel to one side of } \triangle \text{ divides other 2 sides in same ratio}) \\ = \frac{3}{5}$$

$$\therefore \frac{BH}{BC} = \frac{3}{8}$$

In $\triangle GAD$, $\triangle GBH$

\hat{BGH} is common

$$\hat{GAD} = \hat{GBH} \quad (\text{corresp } \angle \text{s equal})$$

$\triangle GAD \sim \triangle GBH$ (equiangular)

$$\therefore \frac{GA}{GB} = \frac{AD}{BH} = \frac{BC}{BH} \quad (BC = AD \text{ opp sides of parallelogram})$$

$$\frac{x+8}{x} = \frac{8}{3}$$

$$3x + 24 = 8x$$

$$24 = 5x$$

$$x = 4.8$$

$$\therefore AG = 12.2 \text{ cm}$$

$$(ii) \text{Area } \triangle CDH = \frac{1}{2} \times CH \times DC \sin \hat{DCH}$$

$$\text{Area } ABCD = BC \times CD \sin \hat{DCH}$$

$$\frac{\text{Area } \triangle CDH}{\text{Area } ABCD} = \frac{\frac{1}{2} \cdot CH \cdot DC \cdot \sin \hat{DCH}}{BC \times DC \cdot \sin \hat{DCH}}$$

$$= \frac{\frac{1}{2} CH}{BC}$$

$$= \frac{1}{2} \times \frac{5}{8}$$

$$= \frac{5}{16}$$

Question 7

$$(a) (i) \frac{d(\frac{1}{2}v^2)}{dx} = 2x^3 - 10x$$

$$\frac{1}{2}v^2 = \frac{2x^4}{4} - \frac{10x^2}{2} + C$$

When $x=0$ $v=u$

$$\frac{1}{2}u^2 = C$$

$$\frac{1}{2}v^2 = \frac{x^4}{2} - \frac{10x^2}{2} + \frac{u^2}{2}$$

$$v^2 - u^2 = x^4 - 10x^2$$

$$(ii) (2) \text{ If } u=3 \text{ then } v^2 - 9 = x^4 - 10x^2$$


$$v^2 = x^4 - 10x^2 + 9$$

$$= (x^2 - 1)(x^2 - 9)$$

$$= (x-1)(x+1)(x-3)(x+3)$$

Since $v^2 \geq 0$ then $(x-1)(x+1)(x-3)(x+3) \geq 0$

$$\therefore x \leq -3 \text{ or } -1 \leq x \leq 1 \text{ or } x \geq 3$$



Since particle starts at $x=0$ with $v=3$ it then moves to the right (positive direction). At $x=1$ $v=0$ and $\dot{x}=-8$ and so particle will then move to the left until it reaches $x=-1$ where $v=0$ and $\dot{x}=8$. This means particle will then move to the right until $v=0$ again at $x=1$. Thus particle oscillates between $x=1$ and $x=-1$ ie within the interval $-1 \leq x \leq 1$.

(B) Not SHM since $\ddot{x} \neq -\omega^2(x-a)$


$$(iii) \text{ If } u=6 \text{ then } v^2 - 36 = x^4 - 10x^2$$

$$v^2 = x^4 - 10x^2 + 25 + 11$$

$$= (x^2 - 5)^2 + 11$$


$\therefore v^2 > 0$ for all x in domain

Particle will never stop



Moves to the right with decreasing velocity until it passes $x=\sqrt{5}$ (with velocity $\sqrt{11}$) and continues to the right with increasing velocity

(4)



$$\begin{aligned}
 y &= 4 - x^2 \\
 y &= 3x \\
 x^2 + 3x - 4 &= 0 \\
 (x+4)(x-1) &= 0 \\
 x &= -4, 1 \\
 y &= -12, 3
 \end{aligned}$$

Let $A(y)$ be the area of the cross-section at height y

$$\begin{aligned}
 A(y) &= \pi R^2 - \pi r^2 & R &= 3 - \frac{y}{3} & y = 3 - \frac{4}{y} \\
 R^2 &= \left(3 - \frac{y}{3}\right)^2 = 9 - 2y + \frac{y^2}{9} & r &= 3 - \sqrt{4-y} & y = 4 - x^2 \\
 r^2 &= \left(3 - \sqrt{4-y}\right)^2 = 9 - 6\sqrt{4-y} + 4-y
 \end{aligned}$$

$$\begin{aligned}
 A(y) &= \pi \left[\left(9 - 2y + \frac{y^2}{9}\right) - \left(9 - 6\sqrt{4-y} + 4-y\right) \right] \\
 &= \pi \left(\frac{y^2}{9} + y + 6\sqrt{4-y} - 4 \right)
 \end{aligned}$$

$$\begin{aligned}
 V &= \int_0^3 \pi \left(\frac{y^2}{9} + y + 6\sqrt{4-y} - 4 \right) dy \\
 &= \pi \left[\frac{y^3}{27} + \frac{y^2}{2} + \frac{6(4-y)^{3/2}}{\frac{3}{2} \times -1} - 4y \right]_0^3 \\
 &= \pi \left\{ \left(\frac{27}{27} - \frac{9}{2} - 4 \times 1 - 12 \right) - \left(-4 \times 4^{3/2} \right) \right\} \\
 &= \pi \left\{ \frac{1}{3} - 16 + 32 \right\} \\
 &= \frac{43\pi}{6}
 \end{aligned}$$

OR cylindrical shells.

Question 8

(a) (i) If x is a factor of both y and z , then $y = ax$ and $z = bx$ where a, b are integers.

$$z - y = bx - ax = (b-a)x \\ \text{i.e. } x \text{ is a factor of } z - y$$

$$(ii) 2^{2^{k+1}} = 2^{2^k \times 2^1} = (2^{2^k})^2$$

$$(iii) F_n = 2^{2^n} + 1 \quad n=0, 1, 2, \dots$$

$$\text{R.T.P } F_n = F_0 F_1 \dots F_{n-1} + 2 \quad n \geq 1$$

$$F_0 = 3$$

$$\begin{aligned} n=1 \quad F_1 &= 2^2 + 1 \\ &= 2^2 + 1 \\ &= 5 \\ &= 3 + 2 \\ &= F_0 + 2 \end{aligned}$$

\therefore Proposition is true for $n=1$

Let k be a value for which proposition is true ($k \geq 1$)

$$\text{i.e. } F_k = F_0 F_1 \dots F_{k-1} + 2$$

Aim to show that proposition is then true for $n=k+1$

$$\text{i.e. } F_{k+1} = F_0 F_1 \dots F_k + 2$$

$$\begin{aligned} F_{k+1} &= 2^{2^{k+1}} + 1 \\ &= (2^{2^k})^2 + 1 \\ &= (F_k - 1)^2 + 1 \\ &= F_k^2 - 2F_k + 1 + 1 \\ &= F_k(F_k - 2) + 2 \\ &= F_k(F_0 F_1 \dots F_{k-1}) + 2 \quad (\text{by inductive hypothesis}) \\ &= F_0 F_1 \dots F_{k-1} F_k + 2 \end{aligned}$$

which is of the required form

\therefore If proposition is true for $n=k$ it is also true for $n=k+1$. Since it

is true for $n=1$ it is also true for $n=1+1=2$ and hence by induction it is true for all positive integers.

(iv) If $m < n$ then

$$F_n = F_0 F_1 \dots F_m F_{m+1} \dots F_{n-1} + 2$$

Let k be the hcf of F_n and F_m .

$\therefore k$ is a factor of F_n and $F_0 F_1 \dots F_m \dots F_{m+1}$

Hence k is a factor of their difference


i.e. k is a factor of 2

i.e. $k = 1$ or 2

But $F_n = 2^{2^n} + 1$ is odd

$\therefore k \neq 2$

i.e. $k = 1$



(i) $F(x) = 0$ when $x = 0, 2, 5+k$ where k is such that $A_1 = A_2$

$$A_1 = \frac{1}{2}(1+3) \times 2 = 4$$

$$A_2 = \frac{1}{2}k^2$$

$$\therefore k^2 = 8$$

$$k = \sqrt{8} = 2\sqrt{2}$$

$$\text{i.e. } x = 0, 2, 5 + 2\sqrt{2}$$

(ii) Stationary points at $x = 1$ and $x = 5$

$$\begin{aligned} F(1) &= \frac{1}{2} \times 1 \times 2 = 1 \\ F(5) &= -4 \end{aligned}$$

$\therefore (1, 1)$ and $(5, -4)$ are stationary points

