



SYDNEY BOYS HIGH SCHOOL
MOORE PARK, SURRY HILLS

2010
HIGHER SCHOOL CERTIFICATE
TRIAL PAPER

Mathematics Extension 2

General Instructions

- Reading Time – 5 Minutes
- Working time – 3 Hours
- Write using black or blue pen. Pencil may be used for diagrams.
- Board approved calculators maybe used.
- Each Question is to be returned in a separate bundle.
- All necessary working should be shown in every question.
- Answer in simplest exact form unless otherwise instructed.

Total Marks – 120

- Attempt questions 1 – 8

Examiner: *A.M.Gainford*

- NOTE: This is a trial paper only and does not necessarily reflect the content or format of the final Higher School Certificate examination paper for this subject.

Question 1. (15 marks) (Start a new answer sheet.)

Marks

2

(a) Evaluate $\int_0^3 \frac{x}{\sqrt{x^2+16}} dx$.

1

(b) Find $\int (\cos^2 x - \sin^2 x) dx$.

2

(c) Use integration by parts to find

$$\int x e^{-x} dx.$$

2

(d) (i) Find real numbers a and b such that

$$\frac{1-3x}{x^2-3x+2} = \frac{a}{x-1} + \frac{b}{x-2}.$$

(ii) Hence find

2

$$\int \frac{1-3x}{x^2-3x+2} dx.$$

2

(e) Evaluate $\int_0^{\frac{\pi}{2}} \frac{1}{\cos x + 2} dx$.

2

(f) (i) If $I_n = \int_0^{\frac{\pi}{4}} \tan^n x dx$, $n = 1, 2, 3, \dots$

show that $I_n + I_{n-2} = \frac{1}{n-1}$, $n = 2, 3, 4, \dots$

2

(ii) Hence evaluate

$$\int_0^{\frac{\pi}{4}} \tan^5 x dx.$$

Question 2. (15 marks) (Start a new answer sheet.)

Marks

(a) If $u = 3 - 4i$ and $v = 2 - 2i$ find **4**

(i) $u\bar{v}$

(ii) \sqrt{u}

(iii) v in modulus-argument form.

(iv) v^4 using De Moivre's theorem.

(b) On an Argand diagram shade the region that is satisfied by both the conditions **2**

$$3 \leq |z - 4i| \leq 4 \text{ and } -\frac{\pi}{4} < \arg(z - 4i) < \frac{\pi}{4}$$

(c) Sketch, on separate Argand diagrams, the locus of the complex number z satisfying **4**

(i) $z^2 - (\bar{z})^2 = i$

(ii) $|z - 1| = \operatorname{Re}(z)$

(d) It is given that $z = \cos \theta + i \sin \theta$ where $0 < \arg z < \frac{\pi}{2}$. **5**

(i) Show that $z + 1 = 2 \cos \frac{\theta}{2} \left(\cos \frac{\theta}{2} + i \sin \frac{\theta}{2} \right)$ and express $z - 1$ in modulus-argument form.

(ii) Hence show that $\operatorname{Re} \left(\frac{z - 1}{z + 1} \right) = 0$.

Question 3. (15 marks) (Start a new answer sheet.)


Marks

- (a) (i) Show that $z = 1 + i$ is a root of $z^2 - (3 - 2i)z + (5 - i) = 0$. **3**
- (ii) Find the other root of the equation.
- (b) If α , β and γ are roots of the equation $x^3 + qx - 2 = 0$ find, in terms of q , the monic cubic polynomial equation whose roots are α^2 , β^2 and γ^2 . **3**
- (c) (i) Use De Moivre's theorem to find $\cos 5\theta$ in terms of powers of $\cos \theta$. **6**
- (ii) Use the result in (i) to solve the equation
- $$16x^4 - 20x^2 + 5 = 0$$
- (d) If ω represents one of the complex roots of the equation $z^3 - 1 = 0$
- (i) Show that $1 + \omega + \omega^2 = 0$.
- (ii) Evaluate $(1 - \omega^8)(1 - \omega^4)(1 - \omega^2)(1 - \omega)$.

Question 4 (15 marks) (Start a new answer sheet.)

(a) The graph of $y = f(x)$ is sketched below.

There is a stationary point at $(0, 1)$.



Use this graph to sketch the following, on separate diagrams, showing essential features.

2

(i) $y = f\left(\frac{x}{2}\right)$

2

(ii) $y = x + f(x)$

2

(iii) $y = \frac{1}{f(x)}$

2

(iv) $y = f\left(\frac{1}{x}\right)$

(b) (i) Find $\int \frac{1}{x^2 \sqrt{9-x^2}} dx$, using the substitution $x = 3\cos\theta$.

4

(ii) Evaluate $\int_1^e x^3 \log_e x dx$.

(c) Given that the quartic polynomial $p(x) = x^4 - 5x^3 - 9x^2 + 81x - 108$ has a zero of multiplicity 3, factorise $p(x)$ completely, and find all its zeroes.

3

Question 5 (15 marks) (Start a new answer sheet.)

Marks

- (a) A particle is moving under gravity in a fluid which exerts a resistance to its motion, per unit mass, k times its speed (k is constant). **2**
- (i) If the particle falls vertically from rest, show that its terminal velocity is $V_T = \frac{g}{k}$, where g is acceleration due to gravity. **2**
- (ii) If the particle is projected vertically upward with velocity V_T show that after time t seconds **6**
- (α) its speed is $V_T(2e^{-kt} - 1)$
- (β) its height above the starting point is $\frac{1}{k}V_T(2 - 2e^{-kt} - kt)$
- (iii) Hence find an expression for the greatest height reached in terms of V_T and k . **2**
- (b) A box contains 6 white balls and 2 black balls. Balls are selected at random, one at a time, and not replaced. A note is kept of the number, X , of the draw which first yields a black ball. If this experiment is repeated many times, find: **5**
- (i) the most probable value of X ;
- (ii) the probability that $X > 4$.


Question 6 (15 marks) (Start a new answer sheet.)

(a) A council has 14 councillors: 6 Labor, 5 Liberal and 3 Independents. Five councillors are chosen at random to form a committee. 6

- (i) (α) How many different committees can be formed?
- (β) Find the probability that the committee will have a majority of Labor councillors.
- (ii) (α) Show that the number of different committees which can be formed with at least one councillor from each of the groups Labor, Liberal, and Independent is 1365.
- (β) Given that the committee contains at least one councillor from each of the groups Labor, Liberal, and Independent, find the probability that the committee will have a majority of Labor councillors.

(b) The circle $x^2 + y^2 = 4$ is rotated about the line $x = 5$ to form a torus. Use the method of cylindrical shells to prove that the volume of the solid is $40\pi^2$ cubic units. 4

(c) The solid drawn at right has a circular base of radius 3 units in the horizontal plane. Vertical cross-sections perpendicular to the diameter along the x -axis are equilateral triangles.



(i) A vertical slice of width Δa is positioned at the point where $x = a$. If the volume of the slice is ΔV , show that $\Delta V = \sqrt{3}(9 - a^2)\Delta a$. 3

(ii) Hence determine the volume of the solid. 2

Question 7 (15 marks) (Start a new answer sheet.)


(a) On polling day in Rock Island City the ratio of electoral votes in the only four polling booths A, B, C, and D was 5:4:3:8 respectively. The percentages of votes for Mr Jones in these booths were 60%, 50%, 40%, and 70% respectively. 4

- (i) Find the probability that a voter chosen at random voted for Mr Jones.
- (ii) If ten voters of this city were chosen at random, find the probability that Mr Jones gained
 - (α) at least 8 votes
 - (β) no more than 2 votes.

(b) The equation $e^{2x} \log_e y = 3$ implicitly defines y as a function of x . 3

Find $\frac{dy}{dx}$ as a function of y .

(c) 8



In the diagram above, P , Q , and R are the midpoints of the sides BC , CA , and AB respectively of a triangle ABC . The circle drawn through the points P , Q , and R meets the sides BC , CA , and AB again at X , Y , and Z respectively.

Copy the diagram to your answer sheet.

- (i) Briefly explain why $RPCQ$ is a parallelogram.
- (ii) Show that ΔXCQ is isosceles.
- (iii) Show that $AX \perp BC$.

Question 8 (15 marks) **(Start a new answer sheet.)**

- (a) Five women and four men are to be seated at a round table. **5**
- (i) In how many ways may this be done without restrictions?
- (ii) In how many ways may this be done if no two men are to be seated together?
- (iii) If one man and one woman are a married couple, what is the probability that they are seated together, given the conditions of part (ii)?

- (b) One root of the equation $x^3 + ax^2 + bx + c = 0$ is equal to the sum of the other two roots. **4**

Show that $a^3 - 4ab + 8c = 0$.

- (c) (i) Graph, in the same xy -plane, the curves **6**
- $y = x^{-\frac{2}{3}}, x > 0$ and $y = (x-1)^{-\frac{2}{3}}, x > 1$
- (ii) Hence, or otherwise, given the sum S , where

$$S = 1 + \frac{1}{\sqrt[3]{2^2}} + \frac{1}{\sqrt[3]{3^2}} + \dots + \frac{1}{\sqrt[3]{(10^9)^2}},$$

find the two consecutive integers between which the sum S lies.

This is the end of the paper.

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, n \neq -1; x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax,$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, a > 0, -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right) x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, x > 0$