



**Question 1** 15 marks

Marks

a) Find  $\int \sin^3 x \, dx$  2

b) Find  $\int \frac{dx}{\sqrt{2x - x^2}}$  2

c) Use partial fractions to find  $\int \frac{5}{(x-3)(2x-1)} \, dx$  3

d) Evaluate  $\int_0^{\frac{\pi}{2}} \frac{dx}{2 + \cos x}$  using the substitution  $t = \tan \frac{x}{2}$  4

e) Find  $\int e^x \sin x \, dx$  4

**Question 2** 15 marks (Use a new page)

Marks

a) Express  $\frac{\overline{2+3i}}{1+i}$  in the form  $x+iy$ . 2

b) Given  $\omega = -1+i\sqrt{3}$  find i)  $|\omega|$  1

ii)  $\arg(\omega)$  1

iii)  $\omega^5 + 16\omega$  in the form  $x+iy$  2

c) On an Argand diagram shade the region specified by

i)  $|z-1| \leq |z-i|$  1

ii)  $\operatorname{Re}\left(\frac{2}{z}\right) \leq 1$  3

d) One of the square roots of  $a+3i$  is equal to  $3+bi$   
where  $a$  and  $b$  are real. Find the value of  $a$  and  $b$ . 2

e) Sketch the region where both the following inequalities hold. 3

$$|z-3| \leq 3 \quad \text{and} \quad 0 \leq \arg(z) \leq \frac{\pi}{2}$$

**Question 3** 15 marks (Use a new page)

Marks

a) Consider the function  $f(x) = \frac{x}{\ln x}$

i) State the natural domain of  $f(x)$  1

ii) Show that the curve  $y = f(x)$  2

has a minimum turning point at the point  $(e, e)$ .

iii) Find the point of inflexion of the curve  $y = f(x)$ . 1

iv) Sketch the curve  $y = f(x)$  showing the important features. 2

b)  $P(cp, \frac{c}{p})$  and  $Q(cq, \frac{c}{q})$  where  $p, q > 0$  are two distinct points on

the hyperbola  $H$  with equation  $xy = c^2$ .

i) Show that the equation of the tangent to  $H$  at  $P(cp, \frac{c}{p})$  2

is given by  $x + p^2y = 2cp$ .

ii) The tangents to  $H$  at  $P(cp, \frac{c}{p})$  and  $Q(cq, \frac{c}{q})$  meet at  $T$ . 2

Find the coordinates of  $T$ .

iii) Find the equation of the chord  $PQ$ . 2

iv) Given that the chord  $PQ$  passes through the point  $(2c, 0)$  1

find a relationship between the parameters  $p$  and  $q$ .

v) Find the equation of the locus of  $T$  as  $P$  and  $Q$  move about  $H$  2

according to the restriction in part iv).

Give a complete description of this locus.

**Question 4** 15 marks (Use a new page)

Marks

- a) The area bounded by  $y = \frac{1}{x+1}$ , the  $x$  axis 4


and the lines  $x = 0$  and  $x = 2$  is rotated about the line  $x = 2$ .

Use cylindrical shells to find the volume of the solid of revolution formed.

- b) Find the equation of the ellipse with centre the origin, 3

which has a focus at  $(2,0)$  and the corresponding directrix is  $x = 4$ .

c)



The diagram shows the graph of the function  $y = f(x)$ .

Draw separate sketches of the following

- i)  $y = f(-x)$  1
- ii)  $y = f|x|$  1
- iii)  $y = \ln(f(x))$  2
- iv)  $x = f(y)$  1
- d) If  $\omega$  is a complex root of  $z^3 = 1$  find the value of


i)  $1 + \omega + \omega^2$  1

ii)  $\frac{1}{3 + 5\omega + 3\omega^2} + \frac{1}{7 + 7\omega + 9\omega^2}$  2

**Question 5** 15 marks (Use a new page)

Marks

a)



Two circles touch internally at  $A$ .  $MT$  is a common tangent.

A tangent to the inner circle at  $P$  cuts the outer circle at  $B$  and  $C$ .

The interval  $AB$  cuts the inner circle at  $Q$ .

The intervals  $PA$ ,  $CA$  and  $PQ$  have been drawn.

Neatly draw the diagram on your answer sheet.

Prove that  $PA$  bisects  $\angle BAC$ .

5

Question 5 (Continued)

Marks

b) The equation  $2x^4 - 3x^2 - 2x + k = 0$  has a triple root.

4

Find the value of  $k$ .

c) Solve  $x^4 - 6x^3 + 15x^2 - 18x + 10 = 0$  over the complex field

4

given that  $1 + i$  is a solution.

d) On an Argand diagram  $\triangle POQ$  is a right angled isosceles triangle.

$O$  is the origin,  $P$  lies in the first quadrant and  $Q$  lies in the second.

$$\angle POQ = 90^\circ.$$

Given that  $OP$  represents the complex number  $a + ib$ ,

Write down the complex number represented by

i)  $OQ$

1

ii)  $PQ$

1

**Question 6** 15 marks (Use a new page)

Marks

a) The function  $f$  is given by


$$f(x) = e^{\frac{x}{1+kx}} \text{ where } k \text{ is a positive constant.}$$

i) Given that the tangent to  $y = f(x)$  at the point  $(a, f(a))$  passes through the origin show that

$$k^2 a^2 + (2k - 1)a + 1 = 0.$$

ii) Deduce that there is no such tangent as in part i) if  $4k > 1$ .

b)



$ABC$  is a triangle. The line  $RQ$  produced meets  $BC$  produced at  $P$ .

$CX$  is drawn parallel to  $QR$ .

Show that  $\frac{BP}{PC} \times \frac{CQ}{QA} \times \frac{AR}{RB} = 1$  4


(This is known as Menelaus' Theorem.)



Question 6 (Continued)

Marks

c)



The base of the above solid is the area enclosed by  $y^2 = 2x$  and  $x = 4$ .

Each cross section of the solid by planes perpendicular to the  $x$  axis is an isosceles triangle with the equal sides meeting above the base of the solid. A typical cross section of thickness  $\Delta x$  has been shaded.

The perpendicular height of the solid at  $x = 4$  is 4 and the perpendicular height of the solid at  $x = 0$  is 3.

A straight line joins the vertex of the triangle at  $x = 4$  to the point 3 above  $x = 0$ .

i) Show that the perpendicular height of the solid as a function of  $x$  2

is given by  $\frac{1}{4}x + 3$

ii) Find the volume of the solid. 4

**Question 7** 15 marks (Use a new page)

Marks

a) Given that  $I_n = \int x^n e^{2x} dx$

i) Show that  $I_n = \frac{x^n e^{2x}}{2} - \frac{n}{2} I_{n-1}$  3

ii) Use the above result to find  $\int x^2 e^{2x} dx$ . 2

b) A particle of mass  $2kg$  is propelled from the origin along the  $x$  axis with an initial velocity of  $Qm/s$ .

The only forces acting on the body in the direction of the  $x$  axis are friction which is a constant  $16$  Newtons and air resistance which equals  $v^2$  Newtons where  $v$  is the velocity of the particle  $t$  seconds after leaving the origin.

i) Explain why  $\frac{dv}{dt} = -8 - \frac{1}{2}v^2$ . 1

ii) Show that  $t = \frac{1}{2} \tan^{-1} \left( \frac{4Q - 4v}{16 + Qv} \right)$  5

iii) By using  $\frac{dv}{dt} = v \frac{dv}{dx}$  4

find an expression for  $v$  in terms of  $x$ .

**Question 8** 15 marks (Use a new page)

Marks

a) Solve the equation

4

$$\sin(x + 10^\circ) = \cos(4x) \quad \text{for } 0^\circ \leq x \leq 180^\circ.$$

b) Use Mathematical Induction to prove that

4

$$\tan \theta + 2 \tan 2\theta + \dots + 2^{n-1} \tan(2^{n-1} \theta) = \cot \theta - 2^n \cot(2^n \theta)$$

for all positive integers  $n$ .

c) The function  $f(x)$  is given, for  $x > 0$ , by

$$f(x) = 2 \log_e x - \frac{x^2 - 1}{x}.$$

i) Show that the only zero of  $f(x)$  occurs at  $x = 1$ .

3

Justify your answer.

ii) Let  $g(x) = \frac{x \log_e x}{x^2 - 1}$ , for  $x > 0$  and  $x \neq 1$ .

4

Show that  $0 < g(x) < \frac{1}{2}$ .

*End of Test.*

## STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

NOTE :  $\ln x = \log_e x, \quad x > 0$